The Multivariable Aleph-function involving the Generalized Mellin-Barnes Contour Integrals

ABDI OLI¹, KELELAW TILAHUN², G. V. REDDY³

^{1,2}*Department of Mathematics,*

Wollo University,

P.O. Box: 1145, Dessie, South Wollo, Amhara Region, Ethiopia.

abdioli30@gmail.com, kta3151@gmail.com

³*Department of Mathematics,*

Jigjiga University,

P.O. Box: 1020, Jigjiga, Ethiopia

gvreddy16673@gmail.com

ABSTRACT

In this paper, we have evaluated three definite integrals involving the product of two hypergeometric functions and multivariable Aleph-function. Certain special cases of the main results are also pointed out.

RESUMEN

En este artículo, hemos evaluado tres integrales definidas que involucran el producto de dos funciones hipergeométricas y la función Aleph multivariada. También se señalan ciertos casos especiales del resultado principal.

Keywords and Phrases: Hypergeometric function, Multivariable Aleph function.

2020 AMS Mathematics Subject Classification: 33C20, 33C05.

1 Introduction

The Aleph-function is among very significant special functions and its closely related ones are widely used in physics and engineering. Therefore they are of high interest to physicists and engineers as well as mathematicians. In recent years, many integral formulas involving a diversity of special functions have been presented by many authors (see e.g., [3, 9, 12, 13, 14, 15, 16]). Motivated by these recent papers, three generalized integral formulae involving product of two hypergeometric functions and multivariable Aleph-function are established in the form of three theorems:

For our study, we recall the following three integral formulas (see [5], p. 77, Equations (3.1), (3.2) and (3.3)):

$$\int_0^\infty \left[\left(\alpha x + \frac{\beta}{x} \right)^2 + \gamma \right]^{-\rho-1} dx = \frac{\sqrt{\pi} \Gamma(\rho + \frac{1}{2})}{2\alpha (4\alpha\beta + \gamma)^{\rho+\frac{1}{2}} \Gamma(\rho + 1)} \quad (1.1)$$

$$\left(\alpha > 0; \beta \geq 0; \gamma + 4\alpha\beta > 0; \Re(\rho) + \frac{1}{2} > 0 \right).$$

$$\int_0^\infty \frac{1}{x^2} \left[\left(\alpha x + \frac{\beta}{x} \right)^2 + \gamma \right]^{-\rho-1} dx = \frac{\sqrt{\pi} \Gamma(\rho + \frac{1}{2})}{2\beta (4\alpha\beta + \gamma)^{\rho+\frac{1}{2}} \Gamma(\rho + 1)} \quad (1.2)$$

$$\left(\alpha \geq 0; \beta > 0; \gamma + 4\alpha\beta > 0; \Re(\rho) + \frac{1}{2} > 0 \right).$$

$$\int_0^\infty \left[\left(\alpha + \frac{\beta}{x^2} \right) \left(\alpha x + \frac{\beta}{x} \right)^2 + \gamma \right]^{-\rho-1} dx = \frac{\sqrt{\pi} \Gamma(\rho + \frac{1}{2})}{(4\alpha\beta + \gamma)^{\rho+\frac{1}{2}} \Gamma(\rho + 1)} \quad (1.3)$$

$$\left(\alpha > 0; \beta \geq 0; \gamma + 4\alpha\beta > 0; \Re(\rho) + \frac{1}{2} > 0 \right).$$

We also recall the following identity involving the hypergeometric series ${}_2F_1(\cdot)$ ([8] p. 75, Theorem 1): If

$$(1-y)^{\alpha+\beta-\gamma} {}_2F_1(2\alpha, 2\beta; 2\gamma; y) = \sum_{k=1}^{\infty} a_k y^k, \quad (1.4)$$

then

$${}_2F_1\left(a, b; c + \frac{1}{2}; y\right) {}_2F_1\left(c-a, c-b; c + \frac{1}{2}; y\right) = \sum_{k=0}^{\infty} \frac{(c)_k}{(c + \frac{1}{2})_k} a_k y^k. \quad (1.5)$$

The multivariable Aleph-function defined by Sharma and Ahmad [6] as:

$$\aleph(z_1, z_2, \dots, z_r) \\ = \aleph_{p_i, q_i, \tau_i; R; p_{i(1)}, q_{i(1)}, \tau_{i(1)}; R^{(1)}, \dots, p_{i(r)}, q_{i(r)}, \tau_{i(r)}; R^{(r)}}^{0, n; m_1 n_1; m_2 n_2; \dots; m_r n_r} \left\{ \begin{array}{c} z_1 \\ \vdots \\ z_r \end{array} \middle| \begin{array}{l} B_1 : B_2 \\ B_3 : B_4 \end{array} \right\}$$

$$= \frac{1}{(2\pi\omega)^r} \int_{L_1} \cdots \int_{L_r} \psi(\varsigma_1, \dots, \varsigma_r) \prod_{i=1}^r (\phi_i(\varsigma_i)(z_i)^{\varsigma_i}) d\varsigma_1 \cdots d\varsigma_r \quad (1.6)$$

where, $\omega = \sqrt{-1}$,

$$B_1 = \left((a_j; \alpha_j^{(1)}, \dots, \alpha_j^{(r)})_{1,n}, \left(\tau_i(a_{ji}; \alpha_{ji}^{(1)}, \dots, \alpha_{ji}^{(r)})_{n+1, p_i} \right) \right)$$

$$B_2 = \left((c_j^{(1)}, \gamma_j^{(r)})_{1, n_1}, \left(\tau_{i(1)}(c_{ji(1)}^{(1)}, \gamma_{ji(1)}^{(1)})_{n_1+1, p_{i(1)}} \right) \right)$$

$$; \cdots; \left((c_j^{(r)}, \gamma_j^{(r)})_{1, n_r}, \left(\tau_{i(r)}(c_{ji(r)}^{(r)}, \gamma_{ji(r)}^{(r)})_{n_r+1, p_{i(r)}} \right) \right)$$

$$B_3 = \left(\tau_i(b_{ji}; \beta_{ji}^{(1)}, \dots, \beta_{ji}^{(r)})_{m+1, q_i} \right)$$

$$B_4 = \left((d_j^{(1)}, \delta_j^{(1)})_{1, m_1}, \left(\tau_{i(1)}(d_{ji(1)}^{(1)}, \delta_{ji(1)}^{(1)})_{m_1+1, q_{i(1)}} \right) \right)$$

$$; \cdots; \left((d_j^{(r)}, \delta_j^{(r)})_{1, m_r}, \left(\tau_{i(r)}(d_{ji(r)}^{(r)}, \delta_{ji(r)}^{(r)})_{m_r+1, q_{i(r)}} \right) \right)$$

and

$$\psi(\varsigma_1, \dots, \varsigma_r) = \frac{\prod_{j=1}^n \Gamma(1 - a_j + \sum_{k=1}^r \alpha_j^{(k)} \varsigma_k)}{\sum_{i=1}^R \left[\tau_i \prod_{j=n+1}^{p_i} \Gamma(a_{ji} - \sum_{k=1}^r \alpha_{ji}^{(k)} \varsigma_k) \prod_{j=1}^{q_i} \Gamma(1 - b_{ji} + \sum_{k=1}^r \beta_{ji}^{(k)} \varsigma_k) \right]}, \quad (1.7)$$

$$\phi_k(\varsigma_k) = \frac{\prod_{j=1}^{m_k} \Gamma(d_j^{(k)} - \delta_j^{(k)} \varsigma_k) \prod_{j=1}^{n_k} \Gamma(1 - c_j^{(k)} + \gamma_j^{(k)} \varsigma_k)}{\sum_{i^{(k)}=1}^{R^{(k)}} \left[\tau_{i^{(k)}} \prod_{j=m_k+1}^{q_{i^{(k)}}} \Gamma(1 - d_{ji^{(k)}}^{(k)} + \delta_{ji^{(k)}}^{(k)} \varsigma_k) \prod_{j=n_k+1}^{p_{i^{(k)}}} \Gamma(c_{ji^{(k)}}^{(k)} - \gamma_{ji^{(k)}}^{(k)} \varsigma_k) \right]}, \quad (1.8)$$

The parameters $d_{ji^{(k)}}^{(k)} (j = m_k + 1, \dots, q_{i^{(k)}}), (k = 1, \dots, r; i = 1, \dots, R \text{ \& } i^{(k)} = 1, \dots, R^{(k)})$ are complex numbers. Also positive real numbers α 's, β 's, γ ' and δ 's for standardization purpose such that

$$U_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} + \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} + \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_{i^{(k)}}} \gamma_{ji^{(k)}}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} - \sum_{j=1}^{m_k} \delta_j^{(k)} - \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_{i^{(k)}}} \delta_{ji^{(k)}}^{(k)} \leq 0 \quad (1.9)$$

The real numbers $\tau_i > 0$ ($i = 1, \dots, R$) and $\tau_{i^{(k)}} > 0$ ($i = 1, \dots, R$). The contour is in the s_k -plane and run from $\sigma - \omega \infty$ to $\sigma + \omega \infty$, where σ is real number with loop, if necessary, ensure that the poles of $\Gamma\left(d_j^{(k)} - \delta_j^{(k)} s_k\right)$ with $j = 1, \dots, m_k$ are separated from those of $\Gamma\left(1 - a_j + \sum_{k=1}^r \alpha_j^{(k)} s_k\right)$ with $j = 1, \dots, n$ and $\Gamma\left(1 - c_j^{(k)} + \gamma_j^{(k)} s_k\right)$ with $j = 1, \dots, n_k$ to the left of the contour L_k . The condition for absolute convergence of multiple Mellin-Barnes type contours (1.6) can be obtained by extension of corresponding conditions for multi variable H-function as: $|\arg z_k| < \frac{1}{2} A_i^{(k)} \pi$ where.

$$A_i^{(k)} = \sum_{j=1}^n \alpha_j^{(k)} - \tau_i \sum_{j=n+1}^{p_i} \alpha_{ji}^{(k)} - \tau_i \sum_{j=1}^{q_i} \beta_{ji}^{(k)} + \sum_{j=1}^{n_k} \gamma_j^{(k)} - \tau_{i^{(k)}} \sum_{j=n_k+1}^{p_{i^{(k)}}} \gamma_{ji^{(k)}}^{(k)} + \sum_{j=1}^{m_k} \delta_j^{(k)} - \tau_{i^{(k)}} \sum_{j=m_k+1}^{q_{i^{(k)}}} \delta_{ji^{(k)}}^{(k)} > 0 \quad (1.10)$$

with $k = 1, \dots, r$; $i = 1, \dots, R$ and $i^{(k)} = 1, \dots, R^{(k)}$.

Remark 1: By setting $\tau_i = \tau_{i^{(k)}} = 1$, the multivariable Aleph function reduces to multivariable I-function (see [4, 7]).

Remark 2: By setting $\tau_i = \tau_{i^{(k)}} = 1$ ($k = 1, \dots, r$) and $R = R^{(1)} = \dots, R^{(r)} = 1$, the multivariable Aleph-function reduces to multivariable H-function defined by Srivastava and Panda [10].

Remark 3: When we set $r = 1$, the multivariable Aleph function reduces to Aleph-function of one variable defined by Sudland [11].

2 Main Results

Theorem 2.1. Let $\alpha > 0$, $\beta \geq 0$, $\gamma + 4\alpha\beta > 0$, $\mu_i > 0$, $\eta \geq 0$, $\Re(\rho) + \frac{1}{2} > 0$; $-\frac{1}{2} < \alpha - \beta - \gamma < \frac{1}{2}$; $\Re\left(\lambda + \mu_i \min_{1 \leq j \leq m_i} \left\{ \frac{\Re(d_j^{(i)})}{\delta_j^{(i)}} \right\}\right) > 0$ ($i = 1, \dots, r$), and $\sigma = \left[\left(\alpha x + \frac{\beta}{x} \right)^2 + \gamma \right]$ then the following formula holds:

$$\begin{aligned} & \int_0^\infty \sigma^{-\rho-1} {}_2F_1\left(\alpha, \beta; \gamma + \frac{1}{2}; \sigma\right) {}_2F_1\left(\gamma - \alpha, \gamma - \beta; \gamma + \frac{1}{2}; \sigma\right) \\ & \quad \times \aleph\left(z_1 \sigma^{-\eta_1}, \dots, z_r \sigma^{-\eta_r}\right) dx \\ & = \frac{\sqrt{\pi}}{2\alpha(4\alpha\beta + \gamma)^{\rho + \frac{1}{2}}} \sum_{h=0}^\infty \frac{1}{(4\alpha\beta + \gamma)^{-h}} \frac{(\gamma)_h}{\left(\gamma + \frac{1}{2}\right)_h} a_h \\ & \quad \times \aleph^{0, n+1: m_1 n_1; m_2 n_2; \dots; m_r n_r}_{p_i+1, q_i+1, \tau_i; R; p_{i^{(1)}}, q_{i^{(1)}}, \tau_{i^{(1)}}; R^{(1)}, \dots, p_{i^{(r)}}, q_{i^{(r)}}, \tau_{i^{(r)}}; R^{(r)}} \end{aligned}$$

$$\left\{ \begin{array}{c} \frac{z_1}{(4\alpha\beta+\gamma)^{\eta_1}} \\ \vdots \\ \frac{z_r}{(4\alpha\beta+\gamma)^{\eta_r}} \end{array} \middle| \begin{array}{c} (-\frac{1}{2}-\rho+h; \eta_1, \dots, \eta_r), B_1 : B_2 \\ (-\rho+h; \eta_1, \dots, \eta_r), B_3 : B_4 \end{array} \right\}. \quad (2.1)$$

Proof. Assume that Ω in L.H.S. of (2.1), then by virtue of equation (1.5) and (1.6), we have the following

$$\begin{aligned} \Omega &= \int_0^\infty \sigma^{-\rho-1} \sum_{h=0}^\infty \frac{(\gamma)_h}{(\gamma+\frac{1}{2})_h} a_h \sigma^h \mathbb{N}_{p_i, q_i, \tau_i; R; p_i(1), q_i(1), \tau_i(1); R^{(1)}, \dots, p_i(r), q_i(r), \tau_i(r); R^{(r)}}^{0, n: m_1 n_1, m_2 n_2, \dots, m_r n_r} \\ &\quad \left\{ \begin{array}{c} z_1 \sigma^{-\eta_1} \\ \vdots \\ z_r \sigma^{-\eta_r} \end{array} \middle| \begin{array}{c} B_1 : B_2 \\ B_3 : B_4 \end{array} \right\} dx \\ &= \int_0^\infty \sigma^{-\rho-1} \sum_{h=0}^\infty \frac{(\gamma)_h}{(\gamma+\frac{1}{2})_h} a_h \sigma^h \\ &\quad \times \left\{ \frac{1}{(2\pi\omega)^r} \int_{L_1} \cdots \int_{L_r} \psi(\varsigma_1, \dots, \varsigma_r) \prod_{i=1}^r (\phi_i(\varsigma_i) (z_i \sigma^{-\eta_i})^{\varsigma_i}) d\varsigma_1 \cdots d\varsigma_r \right\} dx \\ &= \sum_{h=0}^\infty \frac{(\gamma)_h}{(\gamma+\frac{1}{2})_h} a_h \left\{ \frac{1}{(2\pi\omega)^r} \int_{L_1} \cdots \int_{L_r} \psi(\varsigma_1, \dots, \varsigma_r) \prod_{i=1}^r (\phi_i(\varsigma_i) (z_i)^{\varsigma_i}) d\varsigma_1 \cdots d\varsigma_r \right\} \\ &\quad \times \int_0^\infty \sigma^{-\rho-1+h-\sum_{k=1}^s \eta_k \varsigma_k} dx \end{aligned}$$

By using equation (1.1), we can obtain the following equation

$$\begin{aligned} \Omega &= \sum_{h=0}^\infty \frac{(\gamma)_h}{(\gamma+\frac{1}{2})_h} a_h \left\{ \frac{1}{(2\pi\omega)^r} \int_{L_1} \cdots \int_{L_r} \psi(\varsigma_1, \dots, \varsigma_r) \prod_{i=1}^r (\varphi_i(\varsigma_i) (z_i)^{\varsigma_i}) d\varsigma_1 \cdots d\varsigma_r \right\} \\ &\quad \times \frac{\sqrt{\pi} \Gamma(\rho-h+\sum_{k=1}^s \eta_k \varsigma_k + \frac{1}{2})}{2\alpha(4\alpha\beta+\gamma)^{\rho+\frac{1}{2}-h+\sum_{k=1}^s \eta_k \varsigma_k} \Gamma(\rho-h+\sum_{k=1}^s \eta_k \varsigma_k + 1)} \\ &= \sum_{h=0}^\infty \frac{(\gamma)_h}{(\gamma+\frac{1}{2})_h} \frac{a_h}{2\alpha(4\alpha\beta+\gamma)^{\rho+\frac{1}{2}-h+\sum_{k=1}^s \eta_k \varsigma_k} \Gamma(\rho-h+\sum_{k=1}^s \eta_k \varsigma_k + 1)} \\ &\quad \times \frac{1}{(2\pi\omega)^r} \int_{L_1} \cdots \int_{L_r} \psi(\varsigma_1, \dots, \varsigma_r) \prod_{i=1}^r (\varphi_i(\varsigma_i) (z_i)^{\varsigma_i}) d\varsigma_1 \cdots d\varsigma_r \\ &= \frac{\sqrt{\pi}}{2\alpha(4\alpha\beta+\gamma)^{\rho+\frac{1}{2}}} \sum_{h=0}^\infty \frac{(\gamma)_h}{(\gamma+\frac{1}{2})_h} \frac{a_h}{(4\alpha\beta+\gamma)^{-h}} \end{aligned}$$

$$\begin{aligned}
& \times \frac{\Gamma(\rho - h + \sum_{k=1}^s \eta_k \varsigma_k + \frac{1}{2})}{(4\alpha\beta + \gamma)^{\sum_{k=1}^s \eta_k \varsigma_k} \Gamma(\rho - h + \sum_{k=1}^s \eta_k \varsigma_k + 1)} \\
& \times \frac{1}{(2\pi\omega)^r} \int_{L_1} \cdots \int_{L_r} \psi(\varsigma_1, \dots, \varsigma_r) \prod_{i=1}^r (\varphi_i(\varsigma_i)(z_i)^{\varsigma_i}) d\varsigma_1 \cdots d\varsigma_r \\
& = \frac{\sqrt{\pi}}{2\alpha(4\alpha\beta + \gamma)^{\rho + \frac{1}{2}}} \sum_{h=0}^{\infty} \frac{1}{(4\alpha\beta + \gamma)^{-h}} \frac{(\gamma)_h}{(\gamma + \frac{1}{2})_h} a_h x \left\{ \frac{\Gamma(\rho - h + \sum_{k=1}^s \eta_k \varsigma_k + \frac{1}{2})}{\Gamma(\rho - h + \sum_{k=1}^s \eta_k \varsigma_k + 1)} \right\} \\
& \times \frac{1}{(2\pi\omega)^r} \int_{L_1} \cdots \int_{L_r} \psi(\varsigma_1, \dots, \varsigma_r) \prod_{i=1}^r \left(\phi_i(\varsigma_i) \left[\frac{z_i}{(4\alpha\beta + \gamma)^{\eta_i}} \right]^{\varsigma_i} \right) d\varsigma_1 \cdots d\varsigma_r
\end{aligned}$$

we readily arrive at the right hand side of (2.1) in view of the presentation of Aleph function in Mellin Barnes contour integral. $\square$

Theorem 2.2. Let $\alpha \geq 0$, $\beta > 0$, $\gamma + 4\alpha\beta > 0$, $\mu_i > 0$, $\eta \geq 0$, $\Re(\rho) + \frac{1}{2} > 0$; $-\frac{1}{2} < \alpha - \beta - \gamma < \frac{1}{2}$; $\Re\left(\lambda + \mu_i \min_{1 \leq j \leq m_i} \left\{ \frac{\Re(a_j^{(i)})}{\delta_j^{(i)}} \right\}\right) > 0$ ($i = 1, \dots, r$), and $\sigma = \left[\left(\alpha x + \frac{\beta}{x} \right)^2 + \gamma \right]$ then the following formula holds:

$$\begin{aligned}
& \int_0^\infty \frac{1}{x^2} \sigma^{-\rho-1} {}_2F_1\left(\alpha, \beta; \gamma + \frac{1}{2}; \sigma\right) {}_2F_1\left(\gamma - \alpha, \gamma - \beta; \gamma + \frac{1}{2}; \sigma\right) \\
& \quad \times \aleph(z_1 \sigma^{-\eta_1}, \dots, z_r \sigma^{-\eta_r}) dx \\
& = \frac{\sqrt{\pi}}{2\beta(4\alpha\beta + \gamma)^{\rho + \frac{1}{2}}} \sum_{h=0}^{\infty} \frac{1}{(4\alpha\beta + \gamma)^{-h}} \frac{(\gamma)_h}{(\gamma + \frac{1}{2})_h} a_h \\
& \quad \times \aleph^{0, n+1; m_1 n_1; m_2 n_2; \dots; m_r n_r}_{p_i+1, q_i+1, \tau_i; R; p_i(1), q_i(1), \tau_i(1); R^{(1)}, \dots, p_i(r), q_i(r), \tau_i(r); R^{(r)}} \\
& \quad \times \left\{ \begin{array}{c} \frac{z_1}{(4\alpha\beta + \gamma)^{\eta_1}} \\ \vdots \\ \frac{z_r}{(4\alpha\beta + \gamma)^{\eta_r}} \end{array} \middle| \begin{array}{c} (-\frac{1}{2} - \rho + h; \eta_1, \dots, \eta_r), B_1 : B_2 \\ (-\rho + h; \eta_1, \dots, \eta_r), B_3 : B_4 \end{array} \right\}. \tag{2.2}
\end{aligned}$$

Proof. In the similar manner of Theorem 2.1 and using (1.2) we easily arrive at the result (2.2). $\square$

Theorem 2.3. Let $\alpha > 0$, $\beta > 0$, $\gamma + 4\alpha\beta > 0$, $\mu_i > 0$, $\eta \geq 0$, $\Re(\rho) + \frac{1}{2} > 0$; $-\frac{1}{2} < \alpha - \beta - \gamma < \frac{1}{2}$; $\Re\left(\lambda + \mu_i \min_{1 \leq j \leq m_i} \left\{ \frac{\Re(a_j^{(i)})}{\delta_j^{(i)}} \right\}\right) > 0$ ($i = 1, \dots, r$), and $\sigma = \left[\left(\alpha x + \frac{\beta}{x} \right)^2 + \gamma \right]$ then the following formula holds:

$$\begin{aligned}
& \int_0^\infty \left(\alpha + \frac{\beta}{x^2} \right) \sigma^{-\rho-1} {}_2F_1\left(\alpha, \beta; \gamma + \frac{1}{2}; \sigma\right) {}_2F_1\left(\gamma - \alpha, \gamma - \beta; \gamma + \frac{1}{2}; \sigma\right) \\
& \quad \times \aleph(z_1 \sigma^{-\eta_1}, \dots, z_r \sigma^{-\eta_r}) dx
\end{aligned}$$

$$\begin{aligned}
&= \frac{\sqrt{\pi}}{(4\alpha\beta + \gamma)^{\rho + \frac{1}{2}}} \sum_{h=0}^{\infty} \frac{1}{(4\alpha\beta + \gamma)^{-h}} \frac{(\gamma)_h}{\left(\gamma + \frac{1}{2}\right)_h} a_h \\
&\times \mathbb{N}_{p_i+1, q_i+1, \tau_i; R; p_{i(1)}, q_{i(1)}, \tau_{i(1)}; R^{(1)}, \dots, p_{i(r)}, q_{i(r)}, \tau_{i(r)}; R^{(r)}}^{0, n+1: m_1 n_1; m_2 n_2; \dots; m_r n_r} \\
&\left\{ \begin{array}{c} \frac{z_1}{(4\alpha\beta + \gamma)^{\eta_1}} \\ \vdots \\ \frac{z_r}{(4\alpha\beta + \gamma)^{\eta_r}} \end{array} \middle| \begin{array}{c} \left(-\frac{1}{2} - \rho + h; \eta_1, \dots, \eta_r\right), B_1 : B_2 \\ \left(-\rho + h; \eta_1, \dots, \eta_r\right), B_3 : B_4 \end{array} \right\}. \quad (2.3)
\end{aligned}$$

Proof. In the similar way of Theorem 2.1 and using (1.3) we easily arrive at the result (2.3). $\square$

3 Special Cases

- (1) If we put $\tau_i = 1$, in (2.1), (2.2) and (2.3), we get the results in terms of multivariable I-function [4, 7].
- (2) Some suitable parametric changes in (1.1), we obtain single variable I-function, then we arrive at the results due to Chand [1].
- (3) Also, multivariable Aleph function reduces to multivariable H-function with some suitable parameters; we get the known result due to Daiya et al. [2].

4 Conclusion

In this article, we analyze the generalized fractional calculus involving definite integrals of Gradshteyn-Ryzhik of the Multivariable Aleph-function. As the special cases of our main results, which are related to I-function, H-function and G-function, we can also get the number of special functions.

References

- [1] M. Chand, “New Theorems involving the I-function and general class of polynomials”, Global Journal of science Frontier Research, vol. 12, no 4-F, pp. 56–68, 2012.
- [2] J. Daiya, J. Ram and D. Kumar, “The Multivariable H-Function and the General Class of Srivastava Polynomials Involving the Generalized Mellin-Barnes Contour Integrals”, Filomat, vol. 30, no. 6, pp. 1457–1464, 2016.
- [3] D. Kumar, S.D. Purohit and J. Choi, “Generalized fractional integrals involving product of multivariable H-function and a general class of polynomials”, J. Nonlinear Sci. Appl., vol. 9, pp. 8–21, 2016.
- [4] Y. N. Prasad, “Multivariable I-function, Vijnāna Parishad Anusandhan Patrika”, vol. 29, no. 4, pp. 231–235, 1986.
- [5] M. I. Qureshi, K.A. Quraishi and R. Pal, “Some definite integrals of Gradshteyn- Ryzhil and other integrals”, Global Journal of Science Frontier Research, vol. 11, no. 4, pp. 75–80, 2011.
- [6] C. K. Sharma and S.S. Ahmad, “On the multivariable I-function”. Acta ciencia Indica Math, vol. 20, no. 2, pp. 113–116, 1994.
- [7] C. K. Sharma and G.K. Mishra, “Exponential Fourier series for the multivariable I-function”, Acta Ciencia Indica Mathematics, vol. 21, no. 4, pp. 495–501, 1995.
- [8] L. J. Slater, “Generalized Hypergeometric functions”, Cambridge Univ. Press, Cambridge, London and New York, (1996).
- [9] R. K. Saxena, J. Ram and D.L. Suthar, “Unified fractional derivative formulas for the multivariable H-function”, Vijnana Parishad Anusandhan Patrika, vol. 49, no. 2, (2006), pp. 159–175, 2006.
- [10] H. M. Srivastava, R. Panda, “Some expansion theorems and generating relations for the H-function of several complex variables”, Comment. Math. Univ. St. Paul. vol. 24, pp. 119–137, 1975.
- [11] N. Sudland, B. Baumann and T. F. Nonnenmacher, “Open problem: who knows about the Aleph-function?”, Fract. Calc. Appl. Anal., vol. 1, no. 4, pp. 401–402, 1998.
- [12] D. L. Suthar and P. Agarwal, “Generalized Mittag-Leffler function and the multivariable H-function involving the Generalized Mellin-Barnes contour integrals”, Communications in Numerical Analysis, vol. 1, pp. 25–33, 2017.

-
- [13] D. L. Suthar, B. Debalkie, and M. Andualem, “Modified Saigo fractional integral operators involving multivariable H-function and general class of multivariable polynomials”, *Advances in Difference Equation*, vol. 213, 2019. DOI: 10.1186/s13662-019-2150-0
- [14] D. L. Suthar, H. Habenom and H. Tadesse, “Certain integrals involving Aleph function and Wright’s generalized Hypergeometric function”, *Acta Universitatis Apulensis*, vol. 52, pp. 1–10, 2017.
- [15] D. L. Suthar, S. Agarwal and D. Kumar, “Certain integrals involving the product of Gaussian hypergeometric function and Aleph function”. *Honam Mathematical J.*, vol. 41, no. 1, pp. 1–17, 2019.
- [16] D. L. Suthar and B. Debalkie, “A new class of integral relation involving Aleph-functions”, *Surveys in Mathematics and its Applications*, vol. 12, pp. 193–201, 2017.