

RESEARCH ARTICLE

Open Access

# Characterization of *Escherichia coli* virulence genes, pathotypes and antibiotic resistance properties in diarrheic calves in Iran

Masoud Shahrani<sup>1</sup>, Farhad Safarpour Dehkordi<sup>1</sup> and Hassan Momtaz<sup>2\*</sup>

## Abstract

**Background:** Calf diarrhea is a major economic concern in bovine industry all around the world. This study was carried out in order to investigate distribution of virulence genes, pathotypes, serogroups and antibiotic resistance properties of *Escherichia coli* isolated from diarrheic calves.

**Results:** Totally, 76.45% of 824 diarrheic fecal samples collected from Isfahan, Chaharmahal, Fars and Khuzestan provinces, Iran were positive for *E. coli* and all of them were also positive for *cnf2*, *hlyA*, *cdtIII*, *f17c*, *lt*, *st*, *stx1*, *eae*, *ehly*, *stx2* and *cnf1* virulence genes. Chaharmahal had the highest prevalence of STEC (84.61%), while Isfahan had the lowest (71.95%). *E. coli* serogroups had the highest frequency in 1–7 days old calves and winter season. Distribution of ETEC, EHEC, AEEC and NTEC pathotypes among *E. coli* isolates were 28.41%, 5.07%, 29.52% and 3.49%, respectively. Statistical analyses were significant for presence of bacteria between various seasons and ages. All isolates had the high resistance to penicillin (100%), streptomycin (98.25%) and tetracycline (98.09%) antibiotics. The most commonly detected resistance genes were *aadA1*, *sul1*, *aac[3]-IV*, *CITM*, and *dfrA1*. The most prevalent serogroup among STEC was O26.

**Conclusions:** Our findings should raise awareness about antibiotic resistance in diarrheic calves in Iran. Clinicians should exercise caution when prescribing antibiotics.

**Keywords:** Antibiotic resistance profile, Diarrheic calves, *Escherichia coli*, Iran, Serogroups, Virulence genes

## Background

Calf diarrhea is one of the most economic and pervasive concern in veterinary industry all around the world. Infectious agents are the most commonly detected causes of calf diarrhea [1-3]. Several studies have been addressed the high distribution of *Escherichia coli* (*E. coli*) strains in infectious calf diarrhea [3-6]. *Escherichia coli* is a gram-negative, rod-shaped, flagellated, non-sporulating and facultative anaerobic bacterium of the family enterobacteriaceae that classically classified into enterohemorrhagic (EHEC), enterotoxigenic (ETEC), necrotoxicogenic (NTEC), enteroinvasive (EIEC), enteropathogenic (EPEC) and attaching and effacing *E. coli* (AEEC) pathotypes [7]. Intimin genes are present in EPEC and some Shigatoxin-producing *E. coli* (STEC). EPEC strains are defined as *eae*

harbouring diarrhoeagenic *E. coli* that possess the ability to form attaching-effacing (A/E) lesions on intestinal cells and that do not possess Shigatoxin encoding genes [8,9]. Diarrheagenic *Escherichia coli* are now broadly placed into 6 classes based on virulence mechanisms. One of these classes, enterotoxigenic *E. coli*, is the most common cause of diarrhea in beef and dairy calves in the first 4 days of life. Two other diarrheagenic classes, EHEC and EPEC, are important causes of disease in human beings, but less well substantiated causes of diarrhea in calves [8,9]. *E. coli* strains that cause hemorrhagic colitis and hemolytic uremic syndrome in humans, express high levels of Shiga toxin, cause A/E lesions in intestinal epithelial cells, and possess a specific 60-MDa EHEC plasmid are known as EHEC [8,9]. One feature EHEC and EPEC have in common is the causation of intestinal epithelial lesions known as A/E. AEEC is a designation for those *E. coli* strains known to cause A/E lesions or at least carry the genes for this trait, and therefore include organisms that fall into

\* Correspondence: hamomtaz@yahoo.com

<sup>2</sup>Department of Microbiology, College of Veterinary Medicine, Shahrekord Branch, Islamic Azad University, P.O. Box: 166, Shahrekord, Iran  
Full list of author information is available at the end of the article

either the EHEC or EPEC classes. Because cattle are carriers of many different serotypes of EHEC, much emphasis has been placed on the public health and food safety concerns associated with the fecal shedding of these organisms. However, much less emphasis has been given to their roles as diarrheagenic pathogens of cattle [8,9]. In fact, several pathotypes of *E. coli* have been known by presence of certain genes. *EAF* and *bfp* are predominant in EPEC and typical genes of EAEC (*pAA*), *cdt* and *cnf* are the major genes of NTEC and finally *sta*, *stb*, *lt*, *f4*, *f5*, *f18*, *stx1* and *szx2* are the genes of the EHEC pathotype [8,9].

EHEC strains are a subset of STEC strains [3,7]. Infection with STEC strains can result in a spectrum of outcomes, ranging from asymptomatic carriage to uncomplicated diarrhea, bloody diarrhea, hemolytic uremic syndrome (HUS), thrombocytopenia, hemolytic anemia, and acute renal failure [3,7,10]. Most outbreaks and sporadic cases of bloody and non-bloody diarrhea and HUS have been attributed to strains of the STEC serogroups including O157, O26, O103, O111, O145, O45, O91, O113, O121 and O128 [11-15].

Heat-labile enterotoxins (LT) and heat-stable enterotoxins (STa or STb) are two of the most important bacterial virulence factors are able to causes severe diarrhea in calves [3,10]. Also, there are some another virulence factors including phage-encoded cytotoxins, called Shiga toxin 1 (*stx1*) and Shiga toxin 2 (*stx2*), the protein intimin (*eae*) and the plasmid-encoded enterohaemolysin or enterohaemorrhagic *E. coli* haemolysin (*ehly*) which are related to the pathogenesis of STEC strains [16]. The NTEC are able to elaborate two types of cytotoxic necrotizing factors (CNF1 and CNF2). CNF factors are heat-labile proteins which can cause diarrhea. Cytolethal distending toxin (CDT) induces enlargement and death of some cultured eukaryotic cell lines and causing diarrhea. Totally, five different *cdt* alleles (*cdt-I*, *cdt-II*, *cdt-III*, *cdt-IV* and *cdt-V*) have been reported in *E. coli* strains [17-21]. The 31A produces F17c fimbria (formerly called 20 K), which is responsible for *N*-acetyl-D-glucosamine-dependent adhesion of bacteria to calves border villi [5,22]. The F17c includes fimbriae expressed by calf diarrheic [23,24]. Studies in France, Scotland and Belgium showed that pathogenic F17-producing *E. coli* strains represent a significant part of the bacterial strains isolated from diarrheic calves [25].

Diseases caused by *E. coli* often require antimicrobial therapy; however, antibiotic-resistant strains of this bacterium cause longer and more severe illnesses than their antibiotic-susceptible counterparts. Several studies have shown that antibiotic resistance in *E. coli* has increased over time [26-28]. Data on the distribution of serogroups, pathotypes, virulence genes and the antimicrobial resistance properties of *E. coli* strains isolated from diarrheic calves is scarce in Iran. Therefore, the aim of

the present study was to characterize *E. coli* strains isolated from Iranian diarrheic calves at the molecule level and investigate their susceptibility to 13 commonly used antibiotics, as well as investigating seasonal variation in the prevalence and serogroup distribution of *E. coli*.

## Results and discussion

We found that 630 out of 824 samples (76.45%) were positive for *E. coli*. Chaharmahal province had the highest incidence of *E. coli* in diarrhea specimens (84.61%), while Isfahan province had the lowest incidence (71.95%).

Distribution of various pathotypes is shown in Table 1. We found that the incidence of ETEC, EHEC, AEEC and NTEC pathotypes were 28.41%, 5.07%, 29.52% and 3.49%, respectively. In addition, 33.49% diarrheic samples were diagnosed as non detected. We found significant differences between the incidence of ETEC and NTEC ( $P = 0.017$ ), ETEC and EHEC ( $P = 0.023$ ), AEEC and NTEC ( $P = 0.015$ ) and AEEC and EHEC ( $P = 0.021$ ) pathotypes of *E. coli*. *Lt* and *K99* (F5) genes were detected in all of the ETEC pathotype, while the *st* gene

**Table 1 Prevalence of virulence genes in *E. coli* pathotypes isolated from diarrheic calves**

Pathotype	No. positive samples	Virulence gene
Non detected	211 (33.49%)	-
ETEC	179 (28.41%) A, B*	<i>lt</i> , <i>F5</i> : 179 (100%) A, B <i>lt</i> , <i>st</i> , <i>F5</i> : 8(4.46%) a <i>F5</i> , <i>F41</i> : 72 (40.22%) b <i>F5</i> , <i>lt</i> : 179 (100%) <i>F41</i> , <i>lt</i> : 67 (37.43%) <i>F41</i> , <i>st</i> , <i>lt</i> : 4 (2.23%)
EHEC	32 (5.07%) a, c	<i>stx1</i> , <i>eae</i> , <i>ehly</i> : 32 (100%) <i>stx1</i> : 179 (96.23%) A <i>stx2</i> : 106 (56.98%) a <i>eae</i> : 95 (51.07%) a
AEEC	186 (29.52%) C, D	<i>stx1</i> , <i>eae</i> : 91 (48.92%) <i>stx2</i> , <i>eae</i> : 61 (32.79%) <i>stx1</i> , <i>stx2</i> , <i>eae</i> : 34 (18.27%) <i>cnf1</i> , <i>hlyA</i> : 14 (63.63%) <i>cnf2</i> , <i>hlyA</i> : 6 (27.27%) <i>cnf1</i> , <i>cnf2</i> , <i>hlyA</i> : 4 (18.18%)
NTEC	22 (3.49%) b, d	<i>cnf1</i> , <i>cnf2</i> , <i>hlyA</i> , <i>cdtIII</i> : 4 (18.18%) <i>cnf1</i> , <i>hlyA</i> , <i>f17c</i> : 2 (9.09%) <i>cnf2</i> , <i>hlyA</i> , <i>f17c</i> : 2 (9.09%) <i>cnf2</i> , <i>hlyA</i> , <i>cdtIII</i> : 20 (90.90%) <i>cnf2</i> , <i>hlyA</i> , <i>cdtIII</i> , <i>f17c</i> : 5 (22.72%)
Total	630 (76.45%)	

\*Similar capital and small letters in column have a significant differences about  $P < 0.05$ .

was detected in 4.46% of them. *F41* gene was detected in 72 samples of the EHEC pathotype (40.22%), included all *stx1*, *eae*, *ehly* genes, while the distribution of *stx1*, *stx2* and *eae* genes in AEEC pathotype were 96.23%, 56.98% and 51.07%, respectively. In addition, 91 samples (48.92%) had both *stx1* and *eae*, 61 (32.79%) samples had both *stx2* and *eae* and 34 (18.27%) had all *stx1*, *stx2* and *eae* genes. Furthermore, both *cnf1* and *hlyA* genes were detected in 14 (63.63%), both *cnf2* and *hlyA* genes in 6 (27.27%), all *cnf1*, *cnf2* and *hlyA* genes in 4 (18.18%), all *cnf1*, *cnf2*, *hlyA* and *cdtIII* genes in 4 (18.18%), all *cnf1*, *hlyA* and *f17c* genes in 2 (9.09%), all *nf2*, *hlyA* and *f17c* genes in 2 (9.09%), all *cnf2*, *hlyA* and *cdtIII* genes in 20 (90.90%) and finally, all *cnf2*, *hlyA*, *cdtIII* and *f17c* genes in 5 (22.72%) of NTEC pathotypes (Table 1). We also found significant differences between the incidences of *lt* and *st* ( $P = 0.033$ ), *f5* and *f41* ( $P = 0.048$ ), *stx1* and *stx2* ( $P = 0.046$ ) and *stx1* and *eae* ( $P = 0.038$ ) genes.

Seasonal distribution of *E. coli* pathotypes is shown in Table 2. Samples that were collected in the winter had the highest incidence of bacteria (54.92%), while those were collected in summer had the lowest incidence (2.53%). Our results showed significant differences ( $P = 0.018$ ) for presences of *E. coli* strains between cold and warm seasons. The age distribution of the diarrheic calves with regard to infection with *E. coli* is shown in Table 3. We found that the 1–7 day-old calves had the

highest incidence of *E. coli* (39.19%), while the 22–30 day-old calves had the lowest incidence (12.01%). Statistical analysis were significant for the incidence of *E. coli* strains between younger and older calves ( $P = 0.035$  between 1–7 days old and 15–21 and  $P = 0.026$  between 1–7 days old and 22–30).

Antimicrobial resistance of *E. coli* pathotypes isolated from the diarrheic calves is shown in Table 4. Bacterial strains exhibited the highest level of resistance to penicillin (100%), followed by streptomycin (98.25%), tetracycline (98.09%), lincomycin (92.69%) and sulfamethoxazol (90.31%), while STEC and ETEC had the highest resistance profiles overall. Totally, the *E. coli* strains of our investigation had the lowest resistance to nitrofurantoin (23.96%) and cephalothin (52.06%). There were significant differences between resistance of *E. coli* strains to penicillin and nitrofurantoin ( $P = 0.013$ ), penicillin and cephalothin ( $P = 0.045$ ), streptomycin and nitrofurantoin ( $P = 0.015$ ), streptomycin and cephalothin ( $P = 0.050$ ), tetracycline and nitrofurantoin ( $P = 0.018$ ), lincomycin and nitrofurantoin ( $P = 0.026$ ) and sulfamethoxazol and nitrofurantoin ( $P = 0.029$ ).

The distribution of antimicrobial resistance genes within the bacterial pathotypes isolated from diarrheic calves is shown in Table 5. Genes that encode resistance to streptomycin, sulfonamide, gentamicin, ampicillin and trimethoprim antibiotics, i.e., *aadA1*, *sul1*, *aac[3]-IV*,

**Table 2 Prevalence of *E. coli* pathotypes isolated from diarrheic calves in different seasons and provinces**

Province	Season	Non detected	ETEC	STEC	NTEC	No. pos. samples
Isfahan (246)	Autumn (84)	18	7	21	-	46
	Winter (111) A*	39	53	1	9	102
	Spring (35)	7	7	11	1	26
	Summer (16) a	1	-	2	-	3
Chaharmahal (182)	Autumn (61) C	23	9	18	3	53
	Winter (95) B	37	42	11	1	91
	Spring (17)	1	1	-	1	3
	Summer (9) b, c	2	1	4	-	7
Fars (208)	Autumn (84) E	22	6	33	2	63
	Winter (94) D	22	10	35	2	69
	Spring (20)	5	16	1	-	22
	Summer (10) d, e	-	3	-	-	3
Khuzestan (188)	Autumn (63) G	5	1	25	3	34
	Winter (86) F	23	13	48	-	84
	Spring (30)	5	8	8	-	21
	Summer (9) f, g	1	2	-	-	3
Total (824)	Autumn (292) C	68	23	97	8	196
	Winter (386) H	121	118	105	12	346
	Spring (102)	28	32	10	2	72
	Summer (44) h, i	4	6	6	-	16

\*Similar capital and small letters in column have a significant differences about  $P < 0.05$ .

**Table 3 Prevalence of *E. coli* pathotypes isolated from diarrheic calves in different ages and provinces**

Province	Age (day)	Non detected	ETEC	STEC	NTEC	No. pos. samples
Isfahan (246)	1-7 A*	36	42	10	-	88
	8-14	14	6	18	1	39
	15-21 a	1	3	23	2	29
	22-30 a	1	2	14	4	21
Chaharmahal (182)	1-7 B	33	29	9	-	71
	8-14	10	6	18	2	36
	15-21 b	1	2	18	3	24
	22-30 b	1	1	17	4	23
Fars (208)	1-7 C	38	37	7	-	82
	8-14 D	21	5	10	-	36
	15-21 c,d	1	4	17	1	23
	22-30 c,d	2	1	12	1	16
Khozestan (188)	1-7 E	36	36	6	-	78
	8-14	15	3	13	1	32
	15-21 e	1	1	10	1	13
	22-30 e	-	1	16	2	19
Total (824)	1-7 F	143	144	32	-	319
	8-14	60	20	59	4	143
	15-21 f	4	10	68	7	89
	22-30 f	4	5	59	11	79

\*Similar capital and small letters in column have a significant differences about  $P < 0.05$ .

*CITM* and *dfxA1* were the most common antibiotic resistance genes in the diarrheic calves. Interestingly, we found that ETEC and STEC had the highest frequency of antibiotic resistance genes. Statistical analyses were significant between the incidence of *aadA1* and *cmlA* ( $P = 0.018$ ), *aadA1* and *qnr* ( $P = 0.024$ ), *aadA1* and *cat1* ( $P = 0.028$ ), *sul1* and *cmlA* ( $P = 0.019$ ), *sul1* and *cat1* ( $P = 0.029$ ) and *sul1* and *qnr* ( $P = 0.031$ ) antibiotic resistance genes.

Incidence of STEC O-serogroups in diarrheic calves is shown in Table 6. O26 (26.60%) had the highest incidence, followed by O157 (14.67%). In addition, the serogroups of 19 samples (8.71%) cannot be detected. The results of our study showed significant difference between the presence of O26 and O113 ( $P = 0.014$ ), O26 and O121 ( $P = 0.016$ ), O26 and O45 ( $P = 0.020$ ), O26 and O128 ( $P = 0.022$ ), O26 and O145 ( $P = 0.027$ ), O157 and O113 ( $P = 0.018$ ) O157 and O121 ( $P = 0.021$ ) and O157 and O45 ( $P = 0.025$ ) serogroup in calves with diarrhea. The distribution of virulence factors, O-serogroups and antibiotic resistance genes in non-pathogenic *E. coli* strains were lower than those of pathogenic strains. Also, non-pathogenic strains of bacteria were more susceptible to tested antibiotics than pathogenic bacteria.

The high importance of geographical area and season on the incidence of *E. coli* strains in diarrheic calves

were addressed in the present study. Similar results have been reported by Mohamed Ou Said et al. [29]. Season and geography are also known to influence passive transfer of colostral immunoglobulins in calves [30]. Season has a significant effect on the calf mortality [31] as well as on the absorption of immunoglobulins in neonatal calves. One possible explanation for the high prevalence of *E. coli* strains in calves in winter is that climatic variables such as heat, rain and thunderstorms, together with variable barometric pressure may have affected the autonomic nervous systems. These variables could affect immunity, thus making calves more susceptible to infections. Alternatively, the higher prevalence of *E. coli* strains in winter in our study may be related to the fact that the mean serum IgG1 concentrations were low in winter born calves and increased during the spring and summer [32]. The higher mortality rates of 69.6% and 15.36% were observed in winter born calves than 39.4% and 5.97% in summer born calves of Afzal et al. [33] and Sharma et al. [34] investigations. Similar results have been reported previously [35,36].

To the best of our knowledge, this is the first and most comprehensive report on serogroups, pathotypes, virulence genes and antimicrobial resistance properties in *E. coli* strains isolated from diarrheic calves in Iran. Totally, the prevalence of *E. coli* strains isolated from calves with

**Table 4 Antibiotic resistance profile of *E. coli* pathotypes isolated from diarrheic calves (Disk diffusion method)**

	P10*	TE30	S10	C30	SXT	GM10	NFX5	L2	CF30	CIP5	TMP5	F/M300	AM10
Non detected (211)	211	209	210	145	201	179	161	196	102	112	189	32	126
ETEC (179)	179	179	175	126	152	150	101	169	118	120	72	18	116
STEC (218)	218	208	214	192	198	167	123	201	106	146	123	96	202
NTEC (22)	22	22	20	2	8	6	2	18	2	2	8	5	4
Total (630)	630 (100%) A**	618 (98.09%) C	619 (98.25%) B	465 (73.80%)	559 (90.31%) E	502 (79.68%)	387 (61.42%)	584 (92.69%)	328 (52.06%) a,b	380 (60.31%)	392 (62.22%)	151 (23.96%) a,b,c,d,e	448 (71.11%)

\*In this table P10 = penicillin (10 µg/disk); TE30 = tetracycline (30 µg/disk); S10 = streptomycin (10 µg/disk); C30 = chloramphenicol (30 µg/disk); SXT = sulfamethoxazol (25 µg/disk); GM10 = gentamycin (10 µg/disk); NFX5 = enrofloxacin (5 µg/disk); L2 = lincomycin (2 µg/disk); CF30 = cephalothin (30 µg/disk); CIP5 = ciprofloxacin (5 µg/disk); TMP5 = trimethoprim (5 µg/disk); F/M300 = nitrofurantoin (300 µg/disk); AM10 = ampicillin (10 µg/disk).

\*\*Similar capital and small letters in rows have a significant differences about  $P < 0.05$ .

**Table 5 Distribution of antibiotic resistance genes of *E. coli* pathotypes isolated from diarrheic calves in different provinces**

	Province	<i>aadA1</i>	<i>tetA</i>	<i>tetB</i>	<i>dfrA1</i>	<i>qnr</i>	<i>aac (3)-IV</i>	<i>sul1</i>	<i>blaSHV</i>	<i>CITM</i>	<i>cat1</i>	<i>cmIA</i>
Non detected (211)	Isfahan (63)	57	23	16	46	10	61	56	49	54	15	7
	Chaharmahal (59)	58	18	22	49	14	56	59	11	48	10	3
	Fars (44)	38	21	16	38	2	40	42	9	41	-	10
	Khozestan (45)	43	37	37	35	3	13	41	12	43	5	4
	Isfahan (46)	44	36	13	18	5	45	42	28	31	20	7
ETEC (179)	Chaharmahal (36)	34	22	19	17	6	32	29	26	30	26	6
	Fars (44)	39	30	14	14	2	29	32	25	31	13	7
	Khozestan (53)	52	14	17	14	3	36	41	13	9	7	6
SETC (218)	Isfahan (62)	58	30	32	36	32	43	52	20	49	14	-
	Chaharmahal (58)	55	28	26	29	36	49	56	29	52	16	36
	Fars (59)	56	14	16	32	16	46	51	29	55	-	40
	Khozestan (39)	33	26	24	23	17	31	33	8	36	2	18
NTEC (22)	Isfahan (6)	4	2	1	3	2	1	8	1	3	-	1
	Chaharmahal (1) a*	1	-	1	1	-	-	1	-	1	-	-
	Fars (10) A	10	6	-	1	3	1	3	-	1	-	-
Total	Khozestan (5)	2	2	2	1	1	1	2	-	1	-	-
	Isfahan (177) A	163A	91	62	103	49a,b	150	158B	98	137	49a,b	15a,b
	Chaharmahal (154)	148C	68	68	96	56c	137	145	66	131	52c	45
	Fars (157)	143D	71	46	85	23d	116	128	63	128	13d	50
	Khozestan (142) a	130E	79	80	73	24e	81	117	33	89	14e	28

\*Similar capital and small letters in rows have a significant differences about  $P < 0.05$ .

diarrhea in Iran (76.45%, our results) was significantly higher than Egypt (10.36%) [37] and India (42.65%). Mora et al. [38] reported that 12% of the calves and 22% of the farms samples were positive for highly virulent STEC serotype O157:H7. Unfortunately, there were limited investigations related to distribution of *E. coli* bacterium and its pathotypes, serogroups, virulence factors and antibiotic resistance properties.

Another important finding relates to the distributions of several bacterial virulence factors in the diarrheic calves of our investigation. A Brazilian study [39] showed that 49.75% of *E. coli*-positive diarrheic samples was toxigenic and the most prevalent virulence genes were *stx1* (9.7%), *stx2* (6.3%),  $\alpha$ -hemolysin (9.7%), *ehly* (6.8%), *cnf1* (0.5%), *LT-II* (8.3%) and *STa* (3.9%). Nguyen

et al. [3] reported that 31.30% of *E. coli* strains of diarrheic calves had one of the fimbrial genes. They revealed that the incidence of shiga toxin genes, enterotoxins and *eae* gene were 46.29%, 0.92%, and 31.48%, respectively.

Herrera-Luna et al. [40] revealed that 17% of all diarrheic and healthy calves of Australian herds were infected by *E. coli*. They showed that 15.2% of *E. coli* strains harbored the shiga toxin genes including *stx1*, *stx2* and *ehly* and *eae* genes. Low incidence of VTEC phenotype and O157:H7 serotypes of *E. coli* strains of diarrheic calves of Najaf, Iraq were reported by Al-Charrakh and Al-Muhana [41]. Diarrheic calves of Bradford et al. [42] investigation hadn't any *cnf1*, *cnf2*, *stx2*, *stB* and *lt* genes, while the K99 fimbriae, *stA* enterotoxin, *stx1* and *eae* genes were detected in 8, 8, 1 and 1 isolates, respectively which was

**Table 6 Prevalence of STEC serogroups isolated from diarrheic calves in different provinces**

Province	O157	O26	O103	O111	O145	O45	O91	O113	O121	O128	Non detected
Isfahan (62)	16	16	-	3	4	2	8	6	-	4	3
Chaharmahal (58)	2	-	6	7	8	10	10	-	3	6	6
Fars (59)	10	28	9	-	2	-	6	-	-	-	4
Khozestan (39)	4	14	-	6	-	-	2	-	4	3	6
Total (218)	32B*	58A	15	16	14a	12a,b	26	6a,b	7a,b	13a	19

\*Similar capital and small letters in rows have a significant differences about  $P < 0.05$ .


lower than our result. It seems that *stx1*, *stx2*, *eae*, *ehly*, *hlyA*, *lt*, *st*, *cnf1*, *cnf2*, *cdtIII* and *f17c* virulence genes are predominant in *E. coli* strains isolated from calves with diarrhea.

In the present study, 211 *E. coli* strains that were isolated from diarrheic samples hadn't any virulence factors. One possible explanation for this finding is the fact that maybe these strains were non-pathogenic *E. coli* and the animals have diarrhea caused by some other infectious agent. Achá et al. [43] reported that 76% of calves were infected with *E. coli*, while the prevalence of other causative agents including *Salmonella* species and *Campylobacter* species were 2% and 11%, respectively. They also reported that 22/55 (40%) strains from diarrheal calves and 14/88 (16%) strains from healthy calves carried the K99 adhesin ( $P = 0.001$ ).

Our results showed that O26 and O157 were the most common serogroups of our study, while O113 and O121 were the less common. Saridakis et al. [44] showed that O26, O114 and O119 were the most prevalent serogroups in *E. coli* strains isolated from diarrheic calves. Mora et al. [38] showed that 52% of *E. coli* strains

isolated from bovine herds were belonged to O26, O22, O77, O4, O105, O20, O157, O113, and O171 serogroups which was similar to our results.

Bloody diarrhea, non-bloody diarrhea, HUS and other clinical complications of infection with STEC strains are serious among calves, compelling clinicians to consider the provision of early, and empirical antibiotic therapy. However, current recommendations and the available data (although limited in scope and only formally studied for O157 and non-O157-related infections in calves) suggest that antibiotics should be withheld if STEC infection is suspected, given concerns that antibiotics may trigger release of *stx* and progression to diarrhea, resulting in worse clinical outcomes. Furthermore, because inappropriate prescriptions of antibiotics select antibiotic resistance, it was not surprising that our study found that resistance to some antibiotic agents was higher than 80%. The *E. coli* strains of our study were resistant to penicillin (100%), streptomycin (98.25%), tetracycline (98.09%), lincomycin (92.69%), sulfamethoxazol (90.31%), gentamycin (79.68%), chloramphenicol (73.8%), ampicillin (71.11%), trimethoprim (62.22%), enrofloxacin (61.42%)



**Figure 1** Geographical areas of sample collection.

**Table 7 Oligonucleotide primers used for detection of virulence factors, serogroups and antibiotic resistance genes in STEC strains isolated from calves with diarrhea**

Primer name	Sequence (5'-3')	Size of product (bp)	Target gene	References
Hly-F	GAGCGAGCTAAGCAGCTTG	889	<i>hlyEHEC</i>	[6]
Hly-R	CCTGCTCCAGAATAAACCCACA			
Stx1-F	CAGTTAATGTGGTGGCGAAGG	348	<i>Stx1</i>	[53]
Stx1-R	CACCAGACAATGTAACCGCTG			
Stx2-F	ATCCTATTCCTGGGAGTTTACG	584	<i>Stx2</i>	[53]
Stx2-R	GCGTCATCGTATACACAGGAGC			
Eae-F	TGCGGCACAACAGGCGGCGA	629	<i>eae</i>	[6]
Eae-R	CGGTCGCCGCACCAGGATTC			
Cnf1-F	GGGGGAAGTACAGAAGAATTA	1111	<i>cnf1</i>	[21]
Cnf1-R	TTGCCGTCCACTCTCTCACCAGT			
Cnf2-F	TATCATACGGCAGGAGGAAGCACC	1240	<i>cnf2</i>	[21]
Cnf2-R	GTCACAATAGACAATAATTTCCG			
Cdt3-F	GAAAATAAATGGAATATAAATGTCCG	555	<i>cdt-III</i>	[21]
Cdt3-R	TTTGTGTCGGTGCAGCAGGAAAA			
F17c-F	GCAGGAACCGCTCCCTTGGC	416	<i>f17c</i>	[23]
F17c-R	CAACTAACGGGATGTACAGTTTC			
LT-F	GCACACGGAGCTCCTCAGTC	218	<i>ltII</i>	[53]
LT-R	TCCTTCATCCTTTCAATGGCTTT			
ST-F	AGGAACGTACATCATTGCCC	521	<i>st</i>	[54]
ST-R	CAAAGCATGCTCCAGCACTA			
O26-F	CAG AAT GGT TAT GCT ACT GT	423	<i>wzx</i>	[55]
O26-R	CCT ACA TTT GTT TTC GGC ATC			
O103-F	TTGGAGCGTTAACTGGACCT	321	<i>wzx</i>	[55]
O103-R	GCTCCCGAGCACGTATAAG			
O111-F	TAG AGA AAT TAT CAA GTT AGT TCC	406	<i>wzx</i>	[55]
O111-R	ATA GTT ATG AAC ATC TTG TTT AGC			
O145-F	CCATCAACAGATTTAGGAGTG	609	<i>wzx</i>	[55]
O145-R	TTTCTACCGCAATCTATC			
O157-F	CGG ACA TCC ATG TGA TAT GG	259	<i>wzx</i>	[55]
O157-R	TTG CCT ATG TAC AGC TAA TCC			
O45-F	CCG GGT TTC GAT TTG TGA AGG TTG	527	<i>wzx1</i>	[56]
O45-R	CAC AAC AGC CAC TAC TAG GCA GAA			
O91-F	GCTGACCTTCATGATCTGTTGA	291	<i>gnd</i>	[57]
O91-R	TAATTTAACCCGTAGAATCGCTGC			
O113-F	GGGTTAGATGGAGCGCTATTGAGA	771	<i>wzx</i>	[12]
O113-R	AGGTCACCCTCTGAATTATGGCAG			
O121-F	TGGCTAGTGGCATTCTGATG	322	<i>wzx</i>	[58]
O121-R	TGATACTTTAGCCGCCCTTG			
O128-F	GCTTTCTGCCGATATTTGGC		<i>galF</i>	[59]
O128-R	CCGACGGACTGATGCCGGTGATT			
<i>aadA1-F</i>	TATCCAGCTAAGCGCGAACT	447	Streptomycin resistance	[60]
<i>aadA1-R</i>	ATTTGCCGACTACCTTGGTC			
<i>tetA-F</i>	GGTTCACCTCGAACGACGTCA	577	Tetracycline resistance	[60]
<i>tetA-R</i>	CTGTCCGACAAGTTGCATGA			



**Table 7 Oligonucleotide primers used for detection of virulence factors, serogroups and antibiotic resistance genes in STEC strains isolated from calves with diarrhea (Continued)**

<i>tetB</i> -F	CCTCAGCTTCTCAACGCGTG	634	Tetracycline resistance	[60]
<i>tetB</i> -R	GCACCTTGCTGATGACTCTT			
<i>dfrA1</i> -F	GGAGTGCCAAAGGTGAACAGC	367	Trimethoprim resistance	[61]
<i>dfrA1</i> -R	GAGGCGAAGCTTGGGTA AAAAC			
<i>Qnr</i> -F	GGGTATGGATATTATTGATAAAG	670	Fluoroquinolone resistance	[62]
<i>Qnr</i> -R	CTAATCCGGCAGCACTATTTA			
<i>aac</i> [3]-IV-F	CTTCAGGATGGCAAGTTGGT	286	Gentamicin resistance	[63]
<i>aac</i> [3]-IV-R	(TCATCTCGTTCTCCGCTCAT			
<i>Sul1</i> -F	TTCGGCATTCTGAATCTCAC	822	Sulfonamide resistance	[63]
<i>Sul1</i> -R	ATGATCTAACCCCTCGGTCTC			
<i>blaSHV</i> -F	TCGCCTGTGTATTATCTCCC	768	Cephalothin resistance	[63]
<i>blaSHV</i> -R	CGCAGATAAATCACCAATG			
<i>CITM</i> -F	TGGCCAGAACTGACAGGCAA	462	Ampicillin resistance	[63]
<i>CITM</i> -R	TTTCTCCTGAACGTGGCTGGC			
<i>ereA</i> -F	GCCGGTGCTCATGAACCTGAG	419	Erythromycin resistance	[63]
<i>ereA</i> -R	CGACTCTATTGATCAGAGGC			
<i>cat1</i> -F	AGTTGCTCAATGTACTATAACC	547	Chloramphenicol resistance	[63]
<i>cat1</i> -R	TTGTAATTCATTAAGCATTCTGCC			
<i>cmlA</i> -F	CCGCCACGGTGTGTGTATC	698	Chloramphenicol resistance	[63]
<i>cmlA</i> -R	CACCTTGCTGCCATCATTAG			

and ciprofloxacin (60.31%). In terms of antibiotic resistance genes, *aadA1*, *sul1*, *aac*[3]-IV, *CITM* and *dfrA1* were the most commonly detected.

Totally, 73.8% of the STEC strains of our study were resistance to chloramphenicol. Chloramphenicol is a banned antibiotic and the high antibiotic resistance to this drug detected in our study indicates that irregular and unauthorized use of it may have occurred in Iran. Unfortunately, veterinarians in many fields of veterinary such as large animal internal medicine, poultry and even aquaculture, use this antibiotic as a basic one. Therefore, in a short period of time, antibiotic resistance will appear. Previous study showed that in some countries 300,000 kg of antibiotics is used yearly on veterinary prescription in animals [45].

Multidrug resistance of *E. coli* strains have been reported previously [46,47]. Rigobelo et al. [47] reported that the *E. coli* strains had the highest resistance to cephalothin (46.1%), followed by tetracycline (45.7%), trimethoprim-sulfadiazine (43.3%) and ampicillin (41.0%). Multiple resistances of *E. coli* isolates to beta-lactams antibiotics including expanded-spectrum aminoglycosides, cephalosporins, tetracycline sulphonamides, and fluoroquinolones have been reported previously [42]. De Verdier et al. [48] showed that the *E. coli* strains were resistant to sulphonamide, streptomycin, ampicillin and tetracycline. They

reported that 61% of all strains were resistant to more than one antibiotic. Similar investigations have been reported from Sweden [48], United State [49] and Czech [50].

The above data highlight large differences in the prevalence of STEC strains in the different studies, as well as differences in virulence genes and antibiotic resistance properties in the clinical samples. This could be related to differences in the type of samples tested, number of samples, method of sampling, experimental methodology, geographical area, antibiotic prescription preference among clinicians, antibiotic availability, and climate differences in the areas where the samples were collected, which would have differed between each study.

### Conclusion

In conclusion, we identified a large number of pathotypes, serogroups, virulence factors and antibiotic resistance genes and resistance to more than one antibiotic in the *E. coli* strains isolated from diarrheic calves in Iran. Our data indicate that O26 and O157 serogroups are predominant in Iranian diarrheic calves. Marked seasonal, senile and geographical variation was also found. Our data revealed that the O26 serogroup, the *lt*, *f5*, *f41*, *stx1*, *stx2*, *eaeA* and *hlyA* putative virulence genes, the *aadA1*, *sul1*, *aac*[3]-IV, *CITM* and *dfrA1* antibiotic resistance genes, and resistance to penicillin, streptomycin, tetracycline, lincomycin,

**Table 8 PCR programs (temperature and volume) for detection of STEC O-serogroups, virulence factors and antibiotic resistance genes**

Gene	PCR program	PCR volume (50 µL)	
<b>O157, O145, O103, O26, O111</b>	1 cycle: 95°C ----- 3 min.	5 µL PCR buffer 10X	
	30 cycle: 95°C ----- 20 s	1.5 mM MgCl <sub>2</sub>	
	58°C ----- 40 s	200 µM dNTP <sup>e</sup>	
	72°C ----- 30 s	0.5 µM of each primers F & R	
	1 cycle: 72°C ----- 8 min	1.25 U Taq DNA polymerase <sup>e</sup>	
	1 cycle: 94°C ----- 6 min.	2.5 µL DNA template	
	34 cycle: 95°C ----- 50 s	5 µL PCR buffer 10X	
	58°C ----- 70 s	2 mM MgCl <sub>2</sub>	
	72°C ----- 55 s	150 µM dNTP	
	1 cycle: 72°C ----- 10 min	0.75 µM of each primers F & R	
<b>O91, O128, O121, O113, O45</b>	1 cycle: 94°C ----- 6 min.	1.5 U Taq DNA polymerase	
	34 cycle: 95°C ----- 50 s	3 µL DNA template	
	56°C ----- 70 s	5 µL PCR buffer 10X	
	72°C ----- 50 s	1.5 mM MgCl <sub>2</sub>	
	1 cycle: 72°C ----- 6 min	200 µM dNTP	
	1 cycle: 95°C ----- 3 min.	0.5 µM of each primers F & R	
	34 cycle: 94°C ----- 60 s	1 U Taq DNA polymerase	
	56°C ----- 45 s	4 µL DNA template	
	72°C ----- 60 s	5 µL PCR buffer 10X	
	1 cycle: 72°C ----- 10 min	2 mM MgCl <sub>2</sub>	
<b>LT, ST</b>	1 cycle: 94°C ----- 8 min.	2.5 mM MgCl <sub>2</sub>	
	32 cycle: 95°C ----- 60 s	200 µM dNTP	
	55°C ----- 70 s	0.5 µM of each primers F & R	
	72°C ----- 2 min	2 U Taq DNA polymerase	
	1 cycle: 72°C ----- 8 min	3 µL DNA template	
	<b>stx1, stx2, eae, hly, cnf1, cnf2, cdtIII, f17c</b>	1 cycle: 94°C ----- 8 min.	5 µL PCR buffer 10X
		32 cycle: 95°C ----- 60 s	2.5 mM MgCl <sub>2</sub>
		55°C ----- 70 s	200 µM dNTP
		72°C ----- 2 min	0.5 µM of each primers F & R
		1 cycle: 72°C ----- 8 min	2 U Taq DNA polymerase
<b>aadA1, tetA, tetB, dfrA1, qnr, Sul1, aac[3]-IV, blaSHV, CITM, cat1, cmlA</b>		1 cycle: 94°C ----- 8 min.	3 µL DNA template
		32 cycle: 95°C ----- 60 s	5 µL PCR buffer 10X
		55°C ----- 70 s	2.5 mM MgCl <sub>2</sub>
		72°C ----- 2 min	200 µM dNTP
		1 cycle: 72°C ----- 8 min	0.5 µM of each primers F & R

sulfamethoxazol, gentamycin, chloramphenicol, ampicillin, trimethoprim, enrofloxacin and ciprofloxacin were the most commonly detected characteristics of *E. coli* strains

isolated from Iranian diarrheic calves. Hence, judicious use of antibiotics is required by clinicians. We suggested use of disk diffusion method.

## Methods

### Samples and *E. coli* identification

Totally, 824 fecal samples of diarrheic calves were collected randomly during January 2010 to January 2011. Geographical distribution (Isfahan, Chaharmahal, Fars and Khuzestan provinces) (Figure 1), season of samples collection (spring, summer, autumn and winter) and age of diarrheic calves (2 to 30 days) were recorded during sampling. Fecal samples were taken using sterile rectal swabs. All swab samples were placed into tubes containing Stuart medium (Merck, Germany). Samples were immediately transferred to the Microbiology and Infectious Diseases Research Center of the Islamic Azad University of Shahrekord. All samples were diluted in phosphate buffered saline (PBS, Merck, Germany). Then samples were cultured on MacConkey's agar (MC, Merck, Germany) (24 h at 37°C). Lactose positive colonies were cultured on Eosin Methylene Blue (EMB, Merck, Germany) (24 h at 37°C). Metallic green colonies were considered as *E. coli*. Several biochemical tests including Triple Sugar Iron Agar (TSI), Indole, Citrate utilization, Voges-Proskauer, urease and Methyl red tests were used for *E. coli* confirmation. All isolates were stored at -70°C in lactose broth (Merck, Germany) restraining 20% glycerol for further description.

### DNA extraction

All *E. coli* isolates were cultured on Peptone Water (PW, Merck, Germany) (24 h at 37°C). One hundred and fifty microliter of cultured PW media was added to 400 µL sterile distilled water. All components were boiled for 12 min. The mentioned suspension was frozen and then centrifuged at 14,000 rpm for 14 min [51].

### Detection of virulence genes, antibiotic resistance genes and serogroups using Polymerase Chain Reaction (PCR)

The method of Sabat et al. [52] was used in order to *E. coli* confirmation. Several PCR protocols were used to study the presence of virulence genes, serogroups and antibiotic resistance properties of STEC strains. List of primers used for this instance is shown in Table 7. PCR conditions including temperature and volume of each reaction is shown in Table 8. All reactions were performed using the PCR thermocycler (Eppendorf Mastercycler 5330; Eppendorf-Nethel-Hinz GmbH, Hamburg, Germany). Amplified PCR products were electrophoresed on 1.5% agarose gel stained with ethidium bromide. Final products were examined using the gen documentation system. Distilled water and *E. coli* CAPM 5933, CAPM 6006, O159:H20 and O157:K88ac:H19 were used as negative and positive controls, respectively.

### Antibiotic susceptibility testing

The Kirby-Bauer disc diffusion method based on the laboratory protocol of the Clinical and Laboratory

Standards Institute [64] was used for study the antibiotic susceptibility pattern of *E. coli* isolates. All isolates were cultured on Mueller-Hinton agar in an aerobic condition (Merck, Germany) (24 h at 37°C). Susceptibility of *E. coli* isolates to commonly used antibiotic agents was measured and interpreted based on the CLSI protocol. *E. coli* ATCC 25922 was used as quality control.

### Statistical analysis

The SPSS (Statistical Package for the Social Sciences) software (Ver. 16) and Chi-square and Fisher's exact tests were used in order to study the statistical relationship between the incidence of bacterium in various age, season and geographical regions and also between the frequency of various virulence factors, pathotypes, antibiotic resistance genes and serogroups. A *P* value < 0.05 was considered statistically significant.

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

Culture, DNA extraction, PCR techniques and supporting of project was performed by MS and Samples collection, Statistical analysis and writing of manuscript were performed by HM and FSD. All authors read and approved the final manuscript.

### Acknowledgements

These present work was done in the Microbiology laboratory, Islamic Azad University, Shahrekord Branch, Sharekord, Iran. This work was supported by the Islamic Azad University, Shahrekord Branch-Iran. The authors would like to thank Dr. E. Tajbakhsh at the Biotechnology Research Center of the Islamic Azad University of Shahrekord, and Dr. Sh. Nejat Dehkordi at the department of Clinical Sciences, Islamic Azad University of Shahrekord for their important technical and clinical support.

### Author details

<sup>1</sup>Young Researchers and Elites Club, Shahrekord Branch, Islamic Azad University, Shahrekord, Iran. <sup>2</sup>Department of Microbiology, College of Veterinary Medicine, Shahrekord Branch, Islamic Azad University, P.O. Box: 166, Shahrekord, Iran.

Received: 9 January 2014 Accepted: 5 June 2014

Published: 23 June 2014

### References

- Schroeder ME, Bounpheng MA, Rodgers S, Baker RJ, Black W, Naikare H, Velayudhan B, Sneed L, Szonyi B, Clavijo A: Development and performance evaluation of calf diarrhea pathogen nucleic acid purification and detection workflow. *J Vet Diagn Invest* 2012, **24**(5):945-953.
- Sans P, De Fontguyon G: Veal calf industry economics. *Rev Med Vet-Toulouse* 2009, **160**(8-9):420-424.
- Nguyen TD, Vo TT, Vu-Khac H: Virulence factors in *Escherichia coli* isolated from calves with diarrhea in Vietnam. *J Vet Sci* 2011, **12**(2):159-164.
- Wani SA, Hussain I, Nabi A, Fayaz I, Nishikawa Y: Variants of *eae* and *stx* genes of atypical enteropathogenic *Escherichia coli* and non-O157 Shiga toxin-producing *Escherichia coli* from calves. *Lett Appl Microbiol* 2007, **45**(6):610-615.
- Arya G, Roy A, Choudhary V, Yadav MM, Joshi CG: Serogroups, atypical biochemical characters, colicinogeny and antibiotic resistance pattern of Shiga toxin-producing *Escherichia coli* isolated from diarrhoeic calves in Gujarat, India. *Zoonoses Public Health* 2008, **55**(2):89-98.
- Islam MA, Heuvelink AE, de Boer E, Sturm PD, Beumer RR, Zwietering MH, Faruque AS, Haque R, Sack DA, Talukder KA: Shiga toxin-producing *Escherichia coli* isolated from patients with diarrhoea in Bangladesh. *J Med Microbiol* 2007, **56**(Pt 3):380-385.

7. Kaper JB, Nataro JP, Mobley HL: Pathogenic *Escherichia coli*. *Nat Rev Microbiol* 2004, **2**(2):123–140.
8. Moxley RA, Smith DR: Attaching-effacing *Escherichia coli* infections in cattle. *Vet Clin North Am Food Anim Pract* 2010, **26**(1):29–56.
9. Janke BH, Francis DH, Collins JE, Libal MC, Zeman DH, Johnson DD: Attaching and effacing *Escherichia coli* infections in calves, pigs, lambs, and dogs. *J Vet Diagn Invest* 1989, **1**(1):6–11.
10. Kumar A, Taneja N, Singhi S, Sharma RSM: Haemolytic uraemic syndrome in India due to Shiga toxin-producing *Escherichia coli*. *J Med Microbiol* 2012, **62**(pt 1):157–160.
11. Jenkins C, Pearce MC, Smith AW, Knight HI, Shaw DJ, Cheasty T, Foster G, Gunn GJ, Dougan G, Smith HR, Frankel G: Detection of *Escherichia coli* serogroups O26, O103, O111 and O145 from bovine faeces using immunomagnetic separation and PCR/DNA probe techniques. *Lett Appl Microbiol* 2003, **37**(3):207–212.
12. DebRoy C, Roberts E, Kundrat J, Davis MA, Briggs CE, Fratamico PM: Detection of *Escherichia coli* serogroups O26 and O113 by PCR amplification of the wzx and wzy genes. *Appl Environ Microbiol* 2004, **70**(3):1830–1832.
13. Heijnen L, Medema G: Quantitative detection of *E. coli*, *E. coli* O157 and other shiga toxin-producing *E. coli* in water samples using a culture method combined with real-time PCR. *J Water Health* 2006, **4**(4):487–498.
14. Erickson MC, Doyle MP: Food as a vehicle for transmission of Shiga toxin-producing *Escherichia coli*. *J Food Prot* 2007, **70**(10):2426–2449.
15. Lin A, Nguyen L, Lee T, Clotilde LM, Kase JA, Son I, Carter JM, Lauzon CR: Rapid O serogroup identification of the ten most clinically relevant STECs by Luminex microbead-based suspension array. *J Microbiol Methods* 2011, **87**(1):105–110.
16. Law D: Virulence factors of *Escherichia coli* O157 and other Shiga toxin-producing *E. coli*. *J Appl Microbiol* 2000, **88**(5):729–745.
17. Orth D, Grif K, Dierich MP, Würzner R: Cytolethal distending toxins in Shiga toxin-producing *Escherichia coli*: alleles, serotype distribution and biological effects. *J Med Microbiol* 2006, **55**(Pt 11):1487–1492.
18. Bielaszewska M, Stoeve F, Fruth A, Zhang W, Prager R, Brockmeyer J, Mellmann A, Karch H, Friedrich AW: Shiga toxin, cytolethal distending toxin, and hemolysin repertoires in clinical *Escherichia coli* O91 isolates. *J Clin Microbiol* 2009, **47**(7):2061–2066.
19. Tóth I, Nougayrède JP, Dobrindt U, Ledger TN, Boury M, Morabito S, Fujiwara T, Sugai M, Hacker J, Oswald E: Cytolethal distending toxin type I and type IV genes are framed with lambdoid prophage genes in extraintestinal pathogenic *Escherichia coli*. *Infect Immun* 2009, **77**(1):492–500.
20. Janka A, Bielaszewska M, Dobrindt U, Greune L, Schmidt MA, Karch H: Cytolethal distending toxin gene cluster in enterohemorrhagic *Escherichia coli* O157:H- and O157:H7: characterization and evolutionary considerations. *Infect Immun* 2003, **71**(6):3634–3638.
21. Tóth I, Héroult F, Beutin L, Oswald E: Production of cytolethal distending toxins by pathogenic *Escherichia coli* strains isolated from human and animal sources: establishment of the existence of a new cdt variant (Type IV). *J Clin Microbiol* 2003, **41**(9):4285–4291.
22. Ghanbarpour R, Nazem MN: Prevalence of aerobactin and adhesin genes in *Escherichia coli* isolates from blood of bacteremic severely ill neonatal calves. *Vet Arhiv* 2010, **80**(2):185–194.
23. Bertin Y, Girardeau JP, Darfeuille-Michaud A, Martin C: Epidemiological study of pap genes among diarrheagenic or septicemic *Escherichia coli* strains producing CS31A and F17 adhesins and characterization of Pap (31A) fimbriae. *J Clin Microbiol* 2000, **38**(4):1502–1509.
24. Pichon C, Héroult F, du Merle L, Chaudray C, Bonne I, Guadagnini S, Vandewalle A, Le Bouguéne C: Uropathogenic *Escherichia coli* AL511 requires flagellum to enter renal collecting duct cells. *Cell Microbiol* 2009, **11**(4):616–628.
25. De Boer E, Heuvelink AE: Methods for the detection and isolation of Shiga toxin-producing *Escherichia coli*. *Symp Ser Soc Appl Microbiol* 2000, **29**:133S–143S.
26. Solomakos N, Govaris A, Angelidis AS, Pourmaras S, Burriel AR, Kritas SK, Papageorgiou DK: Occurrence, virulence genes and antibiotic resistance of *Escherichia coli* O157 isolated from raw bovine, caprine and ovine milk in Greece. *Food Microbiol* 2009, **26**(8):865–871.
27. Cortés P, Blanc V, Mora A, Dahbi G, Blanco JE, Blanco M, López C, Andreu A, Navarro F, Alonso MP, Bou G, Blanco J, Llagostera M: Isolation and characterization of potentially pathogenic antimicrobial-resistant *Escherichia coli* strains from chicken and pig farms in Spain. *Appl Environ Microbiol* 2010, **76**(9):2799–2805.
28. Tadesse DA, Zhao S, Tong E, Ayers S, Singh A, Bartholomew MJ, McDermott PF: Antimicrobial drug resistance in *Escherichia coli* from humans and food animals, United States, 1950–2002. *Emerg Infect Dis* 2012, **18**(5):742–749.
29. Mohamed Ou Said A, Contrepolis M, Der Vartanian M, Girardeau JP: Factors and markers of virulence of *Escherichia coli* strains isolated from diarrhea in calves aged 4–45 days in Algeria. *Rev Elev Med Vet Pays Trop* 1994, **47**(2):169–175.
30. Snodgrass DR, Terzolo HR, Campbell D, Sherwood I, Menzies JD, Synge BA: Aetiology of diarrhoea in young calves. *Vet Rec* 1986, **119**:31–34.
31. Fink T: Influence of type of housing, microclimate and management on health of calves. Hannover: Inaugural Disser Tierärztliche Hochschule; 1980.
32. Norheim K, Simensen E, Gjestang KE: The relationship between serum IgG levels and age, leg injuries, infections and weight gains in dairy calves. *Nordisk Vet Med* 1985, **37**:113–120.
33. Afzal M, Javed MH, Anjum AD: Calf mortality: seasonal pattern, age distribution and causes of mortality. *Pak Vet J* 1983, **3**:30–33.
34. Sharma MC, Pathak NN, Hung NN, Lien NH, Vuc NV: Mortality in growing Murrah buffalo calves in Vietnam. *Indian J Anim Sci* 1984, **54**:998–1000.
35. Bhullar MS, Tiwana MS: Factors affecting mortality among buffalo calves. *Indian J Anim Sci* 1985, **55**:599–601.
36. Varma AK, Sastry NSR, Kar D: Mortality trends in female Murrah buffalo calves. *Indian J Anim Product Managem* 1988, **4**:18–21.
37. Younis EE, Ahmed AM, El-Khodery SA, Osman SA, El-Naker YF: Molecular screening and risk factors of enterotoxigenic *Escherichia coli* and *Salmonella* spp. in diarrheic neonatal calves in Egypt. *Res Vet Sci* 2009, **87**(3):373–379.
38. Mora A, Herrerra A, López C, Dahbi G, Mamani R, Pita JM, Alonso MP, Llovo J, Bernárdez MI, Blanco JE, Blanco M, Blanco J: Characteristics of the Shiga-toxin-producing enteroaggregative *Escherichia coli* O104:H4 German outbreak strain and of STEC strains isolated in Spain. *Int Microbiol* 2011, **14**(3):121–141.
39. Salvadori MR, Valadares GF, da Silva LD, Blanco J, Yano T: Virulence factors of *Escherichia coli* isolated from calves with diarrhea in Brazil. *Braz J Microbiol* 2003, **34**(3):230–235.
40. Herrera-Luna C, Klein D, Lapan G, Revilla-Fernandez S, Haschek B, Sommerfeld-Stur I, Moestl K, Baumgartner W: Characterization of virulence factors in *Escherichia coli* isolated from diarrheic and healthy calves in Austria shedding various enteropathogenic agents. *Vet Med* 2009, **54**:1–11.
41. Al-Charrakh A, Al-Muhana A: Prevalence of Verotoxin-producing *Escherichia coli* (VTEC) in a survey of dairy cattle in Najaf, Iraq. *Iranian J Microbiol* 2010, **2**:128–134.
42. Bradford PA, Petersen PJ, Fingerling IM, White DG: Characterization of expanded-spectrum cephalosporin resistance in *E. coli* isolates associated with bovine calf diarrhoeal disease. *J Antimicrob Chemother* 1999, **44**(5):607–610.
43. Achá SJ, Kühn I, Jonsson P, Mbazima G, Katouli M, Möllby R: Studies on calf diarrhoea in Mozambique: prevalence of bacterial pathogens. *Acta Vet Scand* 2004, **45**(1–2):27–36.
44. Saridakis HO, el Gared SA, Vidotto MC, Guth BE: Virulence properties of *Escherichia coli* strains belonging to enteropathogenic (EPEC) serogroups isolated from calves with diarrhea. *Vet Microbiol* 1997, **54**(2):145–153.
45. van den Bogaard A: Veterinary use of antibiotics in the Netherlands. Facts and numbers. *Tijdschr Diergeneeskde* 2000, **125**:527–530.
46. Khan A, Das SC, Ramamurthy T, Sikdar A, Khanam J, Yamasaki S, Takeda Y, Nair GB: Antibiotic resistance, virulence gene, and molecular profiles of Shiga toxin-producing *Escherichia coli* isolates from diverse sources in Calcutta, India. *J Clin Microbiol* 2002, **40**(6):2009–2015.
47. Rigobelo EC, Gamez HJ, Marin JM, Macedo C, Ambrosin JA, Ávila FA: Virulence factors of *Escherichia coli* isolated from diarrheic calves. *Arq Bras Med Vet Zootec* 2006, **58**(3):305–310.
48. de Verdier K, Nyman A, Greko C, Bengtsson B: Antimicrobial resistance and virulence factors in *Escherichia coli* from Swedish dairy calves. *Acta Vet Scand* 2012, **54**(1):2.
49. Berge AC, Hancock DD, Sischo WM, Besser TE: Geographic, farm, and animal factors associated with multiple antimicrobial resistance in fecal *Escherichia coli* isolates from cattle in the western United States. *J Am Vet Med Assoc* 2010, **236**(12):1338–1344.

50. Dolejská M, Senk D, Cízek A, Rybáříková J, Sychra O, Literák I: **Antimicrobial resistant *Escherichia coli* isolates in cattle and house sparrows on two Czech dairy farms.** *Res Vet Sci* 2008, **85**(3):491–494.
51. Reischl U, Youssef MT, Kilwinski J, Lehn N, Zhang WL, Karch H, Strockbine NA: **Real-time fluorescence PCR assays for detection and characterization of Shiga toxin, intimin, and enterohemolysin genes from Shiga toxin-producing *Escherichia coli*.** *J Clin Microbiol* 2002, **40**(7):2555–2565.
52. Sabat G, Rose P, Hickey WJ, Harkin JM: **Selective and sensitive method for PCR amplification of *Escherichia coli* 16S rRNA genes in soil.** *Appl Environ Microbiol* 2000, **66**:844–849.
53. Vidal M, Kruger E, Durán C, Lagos R, Levine M, Prado V, Toro C, Vidal R: **Single multiplex PCR assay to identify simultaneously the six categories of diarrheagenic *Escherichia coli* associated with enteric infections.** *J Clin Microbiol* 2005, **43**(10):5362–5365.
54. Arif SK, Salih LIF: **Identification of different categories of diarrheagenic *Escherichia coli* in stool samples by using multiplex PCR technique.** *Asian J Med Sci* 2010, **2**(5):237–243.
55. Possé B, De Zutter L, Heyndrickx M, Herman L: **Metabolic and genetic profiling of clinical O157 and non-O157 Shiga-toxin-producing *Escherichia coli*.** *Res Microbiol* 2007, **158**(7):591–599.
56. DebRoy C, Fratamico PM, Roberts E, Davis MA, Liu Y: **Development of PCR assays targeting genes in O-antigen gene clusters for detection and identification of *Escherichia coli* O45 and O55 serogroups.** *Appl Environ Microbiol* 2005, **71**(8):4919–4924.
57. Perelle S, Dilasser F, Grout J, Fach P: **Identification of the O-antigen biosynthesis genes of *Escherichia coli* O91 and development of a O91 PCR serotyping test.** *J Appl Microbiol* 2002, **93**(5):758–764.
58. Fratamico PM, Briggs CE, Needle D, Chen CY, DebRoy C: **Sequence of the *Escherichia coli* O121 O-antigen gene cluster and detection of enterohemorrhagic *E. coli* O121 by PCR amplification of the *wzx* and *wzy* genes.** *J Clin Microbiol* 2003, **41**(7):3379–3383.
59. Shao J, Li M, Jia Q, Lu Y, Wang PG: **Sequence of *Escherichia coli* O128 antigen biosynthesis cluster and functional identification of an alpha-1,2-fucosyltransferase.** *FEBS Lett* 2003, **553**(1–2):99–103.
60. Randall LP, Cooles SW, Osborn MK, Piddock LJ, Woodward MJ: **Antibiotic resistance genes, integrons and multiple antibiotic resistance in thirty-five serotypes of *Salmonella enterica* isolated from humans and animals in the UK.** *J Antimicrob Chemother* 2004, **53**(2):208–216.
61. Toro CS, Farfán M, Contreras I, Flores O, Navarro N, Mora GC, Prado V: **Genetic analysis of antibiotic-resistance determinants in multidrug-resistant *Shigella* strains isolated from Chilean children.** *Epidemiol Infect* 2005, **133**(1):81–86.
62. Mammeri H, Van De Loo M, Poirel L, Martinez-Martinez L, Nordmann P: **Emergence of plasmid-mediated quinolone resistance in *Escherichia coli* in Europe.** *Antimicrob Agents Chemother* 2005, **49**(1):71–76.
63. Van TT, Chin J, Chapman T, Tran LT, Coloe PJ: **Safety of raw meat and shellfish in Vietnam: an analysis of *Escherichia coli* isolations for antibiotic resistance and virulence genes.** *Int J Food Microbiol* 2008, **124**(3):217–223.
64. CLSI (Clinical and Laboratory Standards Institute): *Performance Standards for Antimicrobial Disk Susceptibility Tests, Approved standard-Ninth Edition (M2-A9)*. Wayne, PA: Clinical and Laboratory Standards Institute; 2006.

doi:10.1186/0717-6287-47-28

**Cite this article as:** Shahrani et al.: Characterization of *Escherichia coli* virulence genes, pathotypes and antibiotic resistance properties in diarrheic calves in Iran. *Biological Research* 2014 **47**:28.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

