

SEPARATION PROBLEM FOR STURM-LIOUVILLE EQUATION WITH OPERATOR COEFFICIENT

Z. OER

Yildiz Technical University, Turkey

Abstract

Let H be a separable Hilbert Space. Denote by $H_1 = L_2(a, b; H)$ the set of function defined on the interval $a < x < b$ ($-\infty \leq a < x < b \leq \infty$) whose values belong to H strongly measurable [12] and satisfying the condition

$$\int_a^b \|f(x)\|_H^2 dx < \infty$$

If the inner product of function $f(x)$ and $g(x)$ belonging to H_1 is defined by

$$(f, g)_1 = \int_a^b (f(x), g(x))_H dx$$

then H_1 forms a separable Hilbert space . We study separation problem for the operator formed by $-y'' + Q(x)y$ Sturm-Liouville differential expression in $L_2(-\infty, \infty; H)$ space has been proved where $Q(x)$ is an operator which transforms at H in value of x , self-adjoint, lower bounded and its inverse is complete continuous.

1. Introduction

Let us consider

$$(1) \quad -y'' + Q(x)y$$

differential expression in $H_1 = L_2(-\infty, \infty; H)$. Let us assume that $Q(x)$, satisfies the following conditions as a self-adjoint operator making transformation at H in each value obtained $(-\infty, \infty)$ interval of x

1) $Q(x)$ operator family have the same definition set indepent of x $(-\infty < x < \infty)$ Let us show this set by \mathcal{D} .

2) Let $Q(x) \geq I$ and for $\forall f \in \mathcal{D}$ $Q(x)f$ be a strong countinuous function in $(-\infty, \infty)$ and $Q^{-1}(x)$ is completely continous in H for $\forall f \in (-\infty, \infty)$

3) When $|x - y| \leq 2$ let

$$\| (Q(x) - Q(y)) Q^{-1}(y) \| < \delta$$

$\delta > 0$ is any number.

Let us form L_0 operator by (2) expression. Let the definition set $\mathcal{D}(L_0)$ of L_0 be by the functions $y(x)$ satisfying the following conditions:

1) Let $y(x)$ having compact support in $(-\infty, \infty)$, $Q(x)y(x), y''(x)$ be continous.

2) Let $-y''(x) + Q(x)y(x) \in L_2(-\infty, \infty; H)$.

We have formed

$$L_0 y = -y'' + Q(x)y \quad , \quad y \in \mathcal{D}(L_0)$$

Since H which is the definition set of $Q(x)$ is dense almost everywhere and since $Q(x)f$ is continuous at $(-\infty, \infty) \forall f \in \mathcal{D}$ definition set $\mathcal{D}(L_0)$ of L_0 operator forms a dense linear monifold almost everywhere of the space $L_2(-\infty, \infty; H)$. L_0 is a symmetric operator bounded from below in $L_2(-\infty, \infty; H)$ Let us assume that L which is a closure of L_0 is a self-adjoint operator.

In this work we study the seperability of the operator L . According to the definition of seperability when $y(x)$ is any function belonging

to $\mathcal{D}(L)$ we will show that $y''(x)$ and $Q(x)y(x)$ functions also belong to $L_2(-\infty, \infty; H)$ space.

Let us show the resolvent of L operator in regular λ value (λ is complex number) by $R_\lambda = (L - \lambda I)^{-1}$. According to the definition of Resolvent, R_λ operator is bounded operator in H_1 space.

Lemma 1 : [9] If $Q(x)R_\lambda$ operator is bounded in H_1 , then L operator is separable in H_1 .

Proof: Let $y(x)$ be an arbitrary element belonging to $\mathcal{D}(L)$ then $(L - \lambda I)y = f$ $f \in H_1$. We can write $y = R_\lambda f$ and $Q(x)y = Q(x)R_\lambda f$ Since $Q(x)R_\lambda$ is a bounded operator, $Q(x)R_\lambda f \in H_1$ i.e $Q(x)y \in H_1$ •

Many books, by B. M. Levitan and I. S. Sargsyan [8], E. C. Titchmarsh [11], M. Otelbayev [10], and papers by T.C.Fulton and S.A.Pruess [6] belonging to singular Sturm-Liouville problem have been written. English mathematicians W. N. Everitt and M. A. Giertz [3], [4], [5] have proved that they introduced separation definition for operator L consisting of expression (1) being real valued function $Q(x)$ with series papers and studies, and separation theorem of $Q(x)$ for operator L in various conditions. For the separability problem, there are works by M. Bayramoglu and A. Abudov [1], K. Boymatov [2], A. Izmaylov and M. Otelbayev [13], M. Otelbayev [14] and many mathematicians; it has taken big place in the book [9] by the K. Minbayev and M. Otelbayev and given many references in the book.

Localization method for separability of L operator. This method was firstly used by R. Ismagilov [7], and were developed by M. Otelbayev [10]. We use that developed method.

Let us show the operator L_j , formed by the following differential expression

$$(2) \quad -y'' + Q(x)y \quad , \quad j - 1 < x < j + 1$$

and

$$(3) \quad y(j - 1) = y(j + 1) = 0 \quad j = 0, \pm 1, \pm 2, \dots \text{ are integers}$$

boundary conditions in space $L_2(j-1, j+1; H)$. Each L_j operator is a positive defined self-adjoint operator.

Let $w(x)$ be a function satisfying the following properties and differentiable defined in $(-\infty, \infty)$

Let

$$w(x) = \begin{cases} 1 & |x| \leq 1 \\ 0 & |x| \geq 1.5 \end{cases}$$

and $\varphi_j(x) = w(x-j)$.

Let us show

$$\psi_j(x) = \begin{cases} \frac{1}{2} & |x-j| \leq 1 \\ 0 & |x-j| > 1 \end{cases}$$

It is seen easily that

$$(4) \quad \sum_{j=-\infty}^{\infty} \psi_j(x) = 1$$

Let $f \in H_1$. $\lambda > 0$ let us show

$$M_\lambda f = \sum_{j=-\infty}^{\infty} \varphi_j (L_j + \lambda I)^{-1} \psi_j f$$

operator by M_λ .

$$(5) \quad (L + \lambda I) M_\lambda f = \sum_{j=-\infty}^{\infty} (L + \lambda I) \varphi_j (L_j + \lambda I)^{-1} \psi_j f$$

Since $L + \lambda I$ and $L_j + \lambda I$ operators coincide in the interval $(j-1.5, j+1.5)$ and $\varphi_j(x)$ has compact support in the interval $(j-1.5, j+1.5)$, we can write

$$(6) \quad \begin{aligned} & (L + \lambda I) M_\lambda f = \\ & - \sum_{j=-\infty}^{\infty} \left[\varphi_j'' (L_j + \lambda I)^{-1} \psi_j f + 2\varphi_j' \frac{d}{dx} (L_j + \lambda I)^{-1} \psi_j f \right] \\ & + \sum_{j=-\infty}^{\infty} \varphi_j (L_j + \lambda I) (L_j + \lambda I)^{-1} \psi_j f \\ & = B_\lambda f + \sum_{j=-\infty}^{\infty} \varphi_j \psi_j f \end{aligned}$$

where

$$B_\lambda f = \sum_{j=-\infty}^{\infty} \left[\varphi_j'' (L_j + \lambda I)^{-1} \psi_j f + 2\varphi_j' \frac{d}{dx} (L_j + \lambda I)^{-1} \psi_j f \right]$$

If we consider $\varphi_j \psi_j = \psi_j$ and condition (4), the expression (6) can be written as

$$(7) \quad (L + \lambda I)M_\lambda f = (I + B_\lambda)f$$

If we apply $(L + \lambda I)^{-1}$ operator to both side of (7) we can write

$$(8) \quad M_\lambda f = (L + \lambda I)^{-1}(I + B_\lambda)f$$

Let $(I + B_\lambda) = g$ or $f = (I + B_\lambda)^{-1}g$. Then (8) equality becomes

$$(9) \quad M_\lambda(I + B_\lambda)^{-1}g = (L + \lambda I)^{-1}g$$

Let us evaluate the norm of B_λ operator transforming in H_1 . Let $f \in H_1$

$$(10) \quad \begin{aligned} \|B_\lambda f\|^2 &= \left\| \sum_{j=-\infty}^{\infty} \left[\varphi_j'' (L_j + \lambda I)^{-1} \psi_j f + 2\varphi_j' \frac{d}{dx} (L_j + \lambda I)^{-1} \psi_j f \right] \right\|^2 \\ &\leq 8 \sum_{j=-\infty}^{\infty} \left\| \varphi_j'' (L_j + \lambda I)^{-1} \psi_j f \right\|^2 \\ &\quad + 8 \sum_{j=-\infty}^{\infty} \left\| \varphi_j' (L_j + \lambda I)^{-1} \psi_j f \right\|^2 \end{aligned}$$

In (10) equality it is considered that support of φ_j' and φ_{j+k} ($k \geq 2$) functions are not intersected. Let us evaluate the terms of the first sum on the right side of (10)

$$(11) \quad \begin{aligned} \left\| \varphi_j'' (L_j + \lambda I)^{-1} \psi_j f \right\|^2 &\leq \frac{c^2}{\lambda^2} \|\psi_j f\|^2 = \frac{c^2}{\lambda^2} \int_{j+1}^{\infty} \|\psi_j f\|_H^2 dx \\ &= \frac{c^2}{\lambda^2} \int_{j-1}^{j+1} \|f(x)\|_H^2 dx \end{aligned}$$

Thus,

$$\left\| \varphi_j'' (L_j + \lambda I)^{-1} \psi_j f \right\|^2 \leq \frac{c^2}{\lambda^2} \int_{j-1}^{j+1} \|f(x)\|_H^2 dx$$

Second sum is

$$\begin{aligned} \left\| \varphi'_j (L_j + \lambda I)^{-1} \psi_j f \right\|^2 &\leq c^2 \left\| \frac{d}{dx} (L_j + \lambda I)^{-1} \right\|^2 \|\psi_j f\|^2 \\ &\leq c^2 \left\| \frac{d}{dx} (L_j + \lambda I)^{-1} \right\|^2 \int_{j-1}^{j+1} \|f(x)\|^2 dx \end{aligned}$$

Let us prove the following lemma.

Lemma 2: $\left\| \frac{d}{dx} (L_j + \lambda I)^{-1} \right\| \leq c \frac{1}{\sqrt{\lambda}}$ ($\lambda > 0$) inequality holds.

Proof: Let us consider $y(j-1) = y(j+1) = 0$ by multiply with $y(x)$ both sides of equation $-y'' + Q(x)y = f$. Then,

$$\begin{aligned} \int_{j-1}^{j+1} (-y'' + Q(x)y, y) dx &= \int_{j-1}^{j+1} (f, y) dx \\ \int_{j-1}^{j+1} [\|y'\|^2 + (Q(x)y, y)] dx &= \int_{j-1}^{j+1} (f, y) dx \end{aligned}$$

If we consider in this inequality that $Q(x) = Q^*(x) \geq I$ and use the Schwartz inequality, we obtain

$$\int_{j-1}^{j+1} \|y'\|^2 \leq \left(\int_{j-1}^{j+1} \|f(x)\|^2 \right)^{1/2} \left(\int_{j-1}^{j+1} \|y(x)\|^2 \right)^{1/2}$$

Since

$$y(x) = (L_j + \lambda I)^{-1} f(x)$$

then

$$\|y(x)\|_{L_2^j}^2 \leq \frac{1}{\lambda^2} \|f\|_{L_2^j}^2 = \frac{1}{\lambda^2} \int_{j-1}^{j+1} \|f(x)\|^2 dx$$

Here $L_2^j = L_2(j-1, j+1)$. If we consider

$$\int_{j-1}^{j+1} \|y'\|^2 dx \leq \frac{1}{\lambda} \int_{j-1}^{j+1} \|f(x)\|^2 dx$$

$$y' = \frac{d}{dx} (L_j + \lambda I)^{-1} f$$

$$\left\| \frac{d}{dx} (L_j + \lambda I)^{-1} f \right\|_{L_2^j} \leq \frac{1}{\sqrt{\lambda}} \|f\|_{L_2^j}$$

is obtain from the last inequality. Since f is the arbitrary element of L_2^j we find

$$\left\| \frac{d}{dx}(L_j + \lambda I)^{-1} \right\| < \frac{1}{\sqrt{\lambda}}$$

Thus, the terms belonging to the second sum in (10) are

$$(12) \quad \left\| \varphi_j' \frac{d}{dx}(L_j + \lambda I)^{-1} \psi_j f \right\|^2 \leq \frac{c}{\lambda} \int_{j-1}^{j+1} \|f(x)\|^2 dx \quad \lambda \geq 1$$

If we consider (10), (11), (12) inequalities we find

$$\begin{aligned} \|B_\lambda f\|^2 &\leq \frac{c^2}{\lambda^2} \sum_{j=-\infty}^{\infty} \int_{j-1}^{j+1} \|f(x)\|^2 dx + \frac{c}{\lambda} \sum_{j=-\infty}^{\infty} \int_{j-1}^{j+1} \|f(x)\|^2 dx \\ &\leq \frac{c_1}{\lambda} \sum_{j=-\infty}^{\infty} \int_{j-1}^{j+1} \|f(x)\|^2 dx \\ &= \frac{c_1}{\lambda} \sum_{j=-\infty}^{\infty} \left(\int_{j-1}^j \|f(x)\|^2 dx + \int_j^{j+1} \|f(x)\|^2 dx \right) \\ &= \frac{c_1}{\lambda} \sum_{j=-\infty}^{\infty} \int_{j-1}^j \|f(x)\|^2 dx + \frac{c_1}{\lambda} \sum_{j=-\infty}^{\infty} \int_j^{j+1} \|f(x)\|^2 dx \\ &= \frac{2c_1}{\lambda} \int_{-\infty}^{\infty} \|f(x)\|^2 dx = \frac{2c_1}{\lambda} \|f\|^2 \end{aligned}$$

Thus,

$$\|B_\lambda\| \leq 2 \frac{c_1}{\lambda}$$

where in big positive values of λ , it is found that $\|B_\lambda\|$ is as small as desired. Therefore, we can write

$$M_\lambda = (L + \lambda I)^{-1}(I + B_\lambda)$$

formula as

$$(L + \lambda I)^{-1} = M_\lambda(I + B_\lambda)^{-1}$$

Lemma 3: If $Q(x)M_\lambda$ is bounded, then L operator is separable.

Proof: According to Lemma 1, if $Q(x)(L + \lambda I)^{-1}$ is bounded, then L is separable. By the equation

$$(L + \lambda I)^{-1} = M_\lambda(I + B_\lambda)^{-1}$$

the proof of Lemma 3 is obtained.

Lemma 4: If $\sup_j \|Q(x)(L_j + \lambda I)^{-1}\| < \infty$, then $Q(x)M_\lambda$ operator is bounded.

Proof:

$$\begin{aligned}
 \|Q(x)M_\lambda f\|^2 &= \left\| \sum_{j=-\infty}^{\infty} \varphi_j(x)(L + \lambda I)^{-1} \psi_j f \right\|^2 \\
 &\leq 4 \sum_{j=-\infty}^{\infty} \left\| \varphi_j Q(x)(L + \lambda I)^{-1} \psi_j f \right\|^2 \\
 &\leq \sum_{j=-\infty}^{\infty} \left\| \varphi_j Q(x)(L + \lambda I)^{-1} \right\|^2 \|\psi_j f\|^2 \\
 &\leq \sup_j \left(\left\| \varphi_j Q(x)(L + \lambda I)^{-1} \right\|^2 \right) \sum_{j=-\infty}^{\infty} \|\psi_j f\|^2 \\
 &= 2A \|f\|_{H_1}^2
 \end{aligned}$$

Thus, $\|Q(x)M_\lambda f\|^2 \leq 2A \|f\|^2$. From this we find

$$\|Q(x)M_\lambda\| \leq \sqrt{2A}$$

and Lemma is proved. Let us show that $\|Q(x)(L_j + \lambda I)^{-1}\|$ is finite. When $|x - y| \leq 2$ let us assume that

$$(13) \quad \|(Q(x) - Q(y))Q^{-1}(y)\| \leq \delta$$

($\delta = \text{const.} > 0$) Let us consider the following boundary value problem

$$-y'' + Q(x)y = f$$

$$y(j-1) = y(j+1) = 0$$

Let us write this problem as

$$(14) \quad -y'' + (-Q(j) + Q(x))y + Q(j)y = f$$

$$y(j-1) = y(j+1) = 0$$

Let

$$-y'' + Q(j)y = v$$

and

$$A_j y = -y'' + Q(j)y$$

Then $v = A_j y$. If we consider these, we can write (14) as

$$\begin{aligned} f = L_j y &= A_j y + (Q(x) - Q(j))y \quad (15) \\ v + (Q(x) - Q(j))A_j^{-1}v &= (I + (Q(x) - Q(j))A_j^{-1})v \quad (1.2) \end{aligned}$$

Let us evaluate the norm of operator $(Q(x) - Q(j))A_j^{-1}$. For this let us write the expression as

$$(Q(x) - Q(j))A_j^{-1} = (Q(x) - Q(j))Q(j)^{-1}(Q(j)A_j^{-1})$$

If we use the condition (13) we can write

$$(16) \quad \|(Q(x) - Q(j))A_j^{-1}\| \leq \delta \|Q(j)A_j^{-1}\|$$

Now let us prove the following Lemma.

Lemma 5: The set of $Q(j)A_j^{-1}$ operator is smooth bounded with 1.

$$\|Q(j)A_j^{-1}\| \leq 1$$

Proof: We will prove the Lemma by using opening formula according to eigenvector of $Q(j)$ operator.

Let us find A_j^{-1} operator in order to show that $\|Q(j)A_j^{-1}\| \leq 1$

Let us consider

$$(17) \quad A_j y = -y'' + Q(j)y = f(x)f(x) \in L_2(j - 1, j + 1; H)$$

$$(18) \quad y(j - 1) = 0 \quad y(j + 1) = 0$$

where $Q^{-1}(j)$ complete continous and therefore its spectrum is pure-disjoint.

Let eigenvalues of $Q(j)$ be

$$\alpha_1(j) \leq \alpha_2(j) \leq \dots$$

and the corresponding eigenvectors to these eigenvalues be

$$g_1(j), g_2(j), \dots$$

These form a base in H .

$$(19) \quad \begin{array}{l} \sum_{k=1}^{\infty} h_k(x) g_k(j) \\ y_j(x) \\ \sum_{k=1}^{\infty} y_{k,j}(x) g_k(j) \end{array} \quad \begin{array}{l} (x) \\ = \\ h_k(x) = (f(x), g_k(j)) \\ = \\ y_{k,j}(x) = (y(x), g_k(j)) \end{array}$$

If we put these equations in (17) we find

$$(20) \quad -\sum_{k=1}^{\infty} y_k''(x) g_k(j) + \sum_{k=1}^{\infty} \alpha_k(j) y_k(x) g_k(j) = \sum_{k=1}^{\infty} h_k(x) g_k(j)$$

$$\begin{array}{l} - y_k''(x) + \alpha_k(j) y_k = h_k(x) \quad k = 1, 2, \dots \\ \sum_{k=1}^{\infty} y_k(x) g_k(j) \Big|_{x=j-1} = 0 \quad , \quad \sum_{k=1}^{\infty} y_k(x) g_k(j) \Big|_{x=j+1} = 0 \\ \sum_{k=1}^{\infty} y_k(j-1) g_k(j) = 0 \quad , \quad \sum_{k=1}^{\infty} y_k(j+1) g_k(j) = 0 \end{array}$$

Thus, (17),(18) problems are transformed

$$(21) \quad -y_k''(x) + \alpha_k(j) y_k = h_k(x)$$

$$(22) \quad y_k(j-1) = 0 \quad , \quad y_k(j+1) = 0$$

The eigenvalues of these problems are

$$\lambda = \frac{k^2\pi^2}{4}$$

The corresponding normalized eigenvectors are the following:

if j is odd;

$$y = \sin \frac{k\pi}{2}x$$

if j even and k is odd;

$$y_{k,j} = \cos \frac{k\pi}{2}x$$

if k is even;

$$y_{k,j} = \sin \frac{k\pi}{2}x$$

$$(23) \quad y_{k,j} = \sum_{m=0}^{\infty} \frac{1}{\lambda_m + \alpha_k(j)} (h_k, \varphi_{m,j}) \varphi_{m,j}$$

$$(h, \varphi_m) = \int_{j-1}^{j+1} h(x) \varphi_m(x) dx$$

Let us consider (23) in (19)

$$y_j(x) = \sum_{k=1}^{\infty} \left(\sum_{m=0}^{\infty} \frac{1}{\lambda_m + \alpha_k(j)} (h_k, \varphi_{m,j}) \varphi_{m,j} \right) g_k(j)$$

$$(24) \quad Q(j)A_j^{-1}f = Q(j)y_j(x)$$

$$\begin{aligned} &= Q(j) \sum_{k=1}^{\infty} \left(\sum_{m=0}^{\infty} \frac{1}{\lambda_m + \alpha_k(j)} (h_k, \varphi_{m,j}) \varphi_{m,j} \right) g_k(j) \\ &= \sum_{k=1}^{\infty} \left(\sum_{m=0}^{\infty} \frac{1}{\lambda_m + \alpha_k(j)} (h_k, \varphi_{m,j}) \varphi_{m,j} \right) Q(j)g_k(j) \\ &= \sum_{k=1}^{\infty} \left[\sum_{m=0}^{\infty} \frac{\alpha_k(j)}{\lambda_m + \alpha_k(j)} (h_k, \varphi_{m,j}) \varphi_{m,j} \right] g_k(j) \end{aligned}$$

Thus,

$$Q(j)f = A_j^{-1}f = \sum_{k=1}^{\infty} \left[\sum_{m=0}^{\infty} \frac{\alpha_k(j)}{\lambda_m + \alpha_k(j)} (h_k, \varphi_{m,j}) \varphi_{m,j} \right] g_k(j)$$

Since $\{\varphi_{m,j}(x)\}_{m=0}^{\infty}$ functions form a complete orthonormal system in $L_2(j-1, j+1)$, and the element $\{g_k(j)\}_{j=1}^{\infty}$ form a complete orthonormal system in H $\{\varphi_{m,j}(x)g_k(j)\}_{m=0, k=1}^{\infty}$ forms a complete orthonormal system in Hilbert space.

Since the system $\{\varphi_{m,j}(x)g_k(j)\}_{m=0, k=1}^{\infty}$ forms a complete orthonormal system in $L_{2,j} = L_2(j-1, j+1; H)$ if we use Parseval equation we can write from (24)

$$(25) \quad \|Q(j)A_j^{-1}f\|^2 = \sum_{k=1}^{\infty} \sum_{m=0}^{\infty} \left(\frac{\alpha_k(j)}{\lambda_m + \alpha_k(j)} \right)^2 |(h_k, \varphi_{m,j})|^2$$

Since $\alpha_k(j) \geq 1$, $\frac{\alpha_k(j)}{\lambda_m + \alpha_k(j)} < 1$ and

$$\|Q(j)A_j^{-1}f\|^2 \leq \sum_{k=1}^{\infty} \sum_{m=0}^{\infty} |(h_k, \varphi_{m,j})|^2 = \|f\|^2$$

or

$$\|Q(j)A_j^{-1}f\| \leq \|f\|$$

that is $\|Q(j)A_j^{-1}\| \leq 1$. Thus, we have proved $Q(j)A_j^{-1}$ operators are straight bounded.

According to Lemma, (16) becomes

$$\|(Q(x) - Q(j))A_j^{-1}\| \leq \delta$$

Let us assume that $\delta < 1$. Then from (15) we find

$$v = (I + (Q(x) - Q(j))A_j^{-1})^{-1}f$$

Let

$$T_j = (I + (Q(x) - Q(j))A_j^{-1})^{-1}$$

The last expression becomes

$$v = T_j f$$

and we obtain

$$y(x) = A_j^{-1}T_j f$$

Thus

$$(26) \quad \begin{aligned} \|Q(x)y(x)\| &= \|(Q(x)Q^{-1}(j))(Q(j)A_j^{-1}T_j f)\| \\ &\leq \|Q(x)Q^{-1}(j)\|c\|T_j f\| \\ &\leq \|Q(x)Q^{-1}(j)\|c_1\|f\|_{L_{2,j}} \end{aligned}$$

If we consider here condition

$$\|(Q(x) - Q(j))Q^{-1}(j)\| \leq \delta$$

then we find

$$\|Q(x)Q^{-1}(j) - I\| \leq \delta$$

Therefore, inequality (26) becomes

$$\|Q(x)y(x)\| \leq c_2\|f\|_{L_{2,j}}$$

Here, if we consider $y(x) = L_j^{-1}f$, the last inequality takes the form of

$$\|Q(x)L_j^{-1}f\| \leq c_2\|f\|_{L_{2,j}}$$

If we consider, then we find

$$\|Q(x)(L_j + \lambda I)^{-1}f\| \leq c_2\|f\|_{L_{2,j}}$$

or

$$\|Q(x)(L_j + \lambda I)^{-1}\| \leq c$$

Thus, we have proved the theorem of separability.

Theorem: When the conditions 1) - 3) are satisfied, L operator is separable in H_1 .

Acknowledgement

I express deep gratitude to M.Bayramoglu and M.Otelbayev for their attention to the work

References

- [1] Bayramoglu, M., Abudov, A. A., *About Sturm-Liouville Operator With Operator Coefficient is Essential Self-Adjoint*, *Sbornik, Spectral Theory of Operators*, Bak, "Elm", pp. 12-20, (1982).
- [2] Boymatov, K. Ch., *Separable Theorems For Sturm-Liouville Operator*, *Matem. Zametki*, Vol.14, No:3, pp. 349-359, (1973).
- [3] Everitt, W. N., Giertz, M. A., *Dirichlet Type Result For Ordinary Differential Operators*, *Math. Ann.*, Vol. 203, No 2, pp. 119-128, (1973).
- [4] Everitt, W.N., Giertz, M.A., *An Example Concerning The Separation Property For Differential Operators*, *Proc.Roy.Soc. Edinburg A*, Vol.71, pp. 159-165, (1973).
- [5] Everitt, W.N., Giertz, M.A., *Inequalities And Separation For Schrödinger Type Operators in $L_2(R_n)$* , *Proc.Roy.Soc. Edinburg A*, Vol.79, pp. 257-265, (1977).
- [6] Fulton, T. C., Pruess, S. A., *Eigenvalue And Eigenfunction Asymptotics For Regular Sturm-Liouville Problems*, *J. Math. Anal. Appl.*, 188, pp. 297-340, (1994).
- [7] Ismagilov, R. S., *Self-adjointness of the Sturm-Liouville Operator* *UMN*, 18, No 5, pp. 161-166, (1963).
- [8] Levitan, B. M. and Sargsyan, I. S., *Sturm-Liouville And Dirac Operators*, Kluzer, Dordrechz, (1991).
- [9] Minbayev, K. T., Otelbayev, M. O., *The Weighted of Functional Spaces and Spectrum of the Differential Operators*, Moskova "Nauka", (1988) (R).
- [10] Otelbayev, M. O., *Behaviour of Spectrum of the Sturm-Liouville operator*, Alma Ata. Bilim (1990).

- [11] Titchmarsh, E. C., *Eigenfunctions Expansions Associated With Second Order Differential Equations*, 2nd ed., Vol.I, Oxford Univ.Press, London, (1962).
- [12] Yosida, K., *Functional Analysis*, Berlin-Gttingen-Heidelberg: Springer-Verlag, (1980).
- [13] Izmaylov, A. L., Otelbayev, M. O., *Weighted Integrability of Solutions of Differential Equation Given in Bounded Area*, Izv. An Kaz. SSR., Ser. Fiz. Mat, No 1, pp. 36-40, (1977).
- [14] Otelbayev, M. O., *Weighted Integrability of Solutions Of Sturm-Liouville Equation*, Matem. Zametki, Vol. 16, No 6, pp. 969-980, (1974).

Received : July 2000.

Z. OER

Department of Mathematics

Faculty of Art and Science

Yildiz Technical University

34210, Davutpasa

Istanbul

Turkey

E-MAIL : zoer@yildiz.edu.tr