

Proyecciones  
Vol. 25, N° 3, pp. 261-271, December 2006.  
Universidad Católica del Norte  
Antofagasta - Chile

## ON SOME FUNCTIONS CONCERNING FUZZY PG-CLOSED SETS

M. CALDAS  
UNIVERSIDADE FEDERAL FLUMINENSE, BRASIL

G. NAVALAGI  
KLE SOCIETYS, INDIA

R. SARAF  
GOVT. K. N. G. COLLEGE, INDIA

Received : May 2006. Accepted : October 2006

### Abstract

In this paper we consider new weak and strong forms of fuzzy preirresoluteness and fuzzy pre-closureness via the concept of fuzzy pg-closed sets which we call ap-Fp-irresolute functions, ap-Fp-closed functions and contra fuzzy pre-irresolute functions and we use it to obtain a characterization of fuzzy pre- $T_{1/2}$  spaces.

2000 Math. Subject Classification : Primary: 54A40

Key Words and Phrases : Fuzzy topological spaces, Fpg-closed sets, fuzzy preopen sets, fuzzy preclosed functions, fuzzy pre-irresolute functions, fuzzy pre- $T_{1/2}$  spaces

## 1. Introduction and Preliminaries

In 1991 A. S. Binshahan [4] introduced and investigated the notions of fuzzy preopen and fuzzy preclosed sets. Since then the advent of these notions, several research papers with interesting results in different respects came to existence ([1],[3],[5],[6],[7],[10],[12]).

In this paper we shall introduce new generalizations of fuzzy pre-closedness called ap-fuzzy pre-closedness by using fuzzy pg-closed sets and study some of their basic properties. This definition enables us to obtain conditions under which functions and inverse functions preserve fuzzy pg-closed sets, also in this paper we present a new generalization of fuzzy pre-irresoluteness called contra fuzzy pre-irresolute. We define this last class of functions by the requirement that the inverse image of each fuzzy pre-open set in the co-domain is fuzzy preclosed in the domain. This notion is a stronger form of ap-fuzzy pre-irresolute and ap-fuzzy preclosed functions. Finally, we also characterize the class of fuzzy pre- $T_{1/2}$  spaces in terms of ap-fuzzy pre-irresolute and ap-fuzzy preclosed functions.

Throughout this paper  $(X, \tau)$ ,  $(Y, \sigma)$  and  $(Z, \gamma)$  (or simply  $X, Y$  and  $Z$ ) represent non-empty fuzzy topological spaces on which no separation axioms are assumed, unless otherwise mentioned. Let  $A$  be a fuzzy subset of  $X$  then  $Cl(A)$  and  $Int(A)$  denote the closure of  $A$  and the interior of  $A$ , respectively.

**1.1 DEFINITION :** The fuzzy set  $A$  of  $X$  is called fuzzy preopen [4] if  $A \cdot IntCl(A)$ .

The fuzzy subset  $B$  of  $X$  is called fuzzy preclosed if, its complement  $B^c$  is fuzzy preopen in  $X$ .

By  $FPO(X)$  (rep.  $FPC(X)$ ), we denote the family of all fuzzy preopen (resp. fuzzy preclosed) sets of  $X$ .

**1.2 DEFINITION :** The intersection of all fuzzy preclosed sets containing  $A$  is called the preclosure of  $A$  and is denoted by  $pCl(A)$  [4].

**1.3 DEFINITION :** The fuzzy interior of  $A$  [4] is denoted by  $plnt(A)$ , and is defined by the union of all fuzzy preopen sets of  $X$  which are contained in  $A$ .

**1.4 DEFINITION :** Recall that a fuzzy set  $A$  is called fuzzy regular open (resp. regular closed) if  $A = IntCl(A)$  (resp.  $A = ClInt(A)$ )[2].

**1.5 DEFINITION :** A function  $f : (X, \tau) \rightarrow (Y, \sigma)$  is said to be :

(i) fuzzy preirresolute [8] if,  $f^{-1}(V)$  is fuzzy preopen set in  $(X, \tau)$  for each fuzzy open set  $V$  of  $(Y, \sigma)$ ,

(ii) fuzzy M-preclosed [11] (resp. fuzzy M-preopen[11]) if, for every fuzzy preclosed (resp. fuzzy preopen) set  $B$  of  $(X, \tau)$ ,  $f(B)$  is fuzzy preclosed (resp. fuzzy preopen) set in  $(Y, \sigma)$ ,

(iii) fuzzy contra preclosed if  $f(A)$  is fuzzy preopen set in  $(Y, \sigma)$  for every fuzzy closed set  $A$  of  $(X, \tau)$ [9].

## 2. ap-Fp-IRRESOLUTE AND ap-Fp-CLOSED FUNCTIONS

**2.1 DEFINITION :** A fuzzy set  $A$  of  $(X, \tau)$  is said to be fuzzy pre-generalized closed (briefly, Fpg-closed) if,  $pCl(A) \subseteq H$  whenever  $A \subseteq H$  and  $H$  is fuzzy preopen in  $X$ . A fuzzy set  $B$  is said to be fuzzy pre-generalized open (briefly, Fpg-open) in  $X$  if its complement  $B^c$  is Fpg-closed set in  $X$ .

**2.2 REMARK :** (i) Every fuzzy preclosed set is Fpg-closed but not conversely.

(ii) The concept of fuzzy g-closed sets and Fpg-closed sets are independent. For,

**2.3 EXAMPLE :** Let  $X = \{a, b\}$ , fuzzy sets  $A, B, E, H$  and  $K$  are defined as :

$$A(a) = 0.3, A(b) = 0.4 ;$$

$$B(a) = 0.6, B(b) = 0.5 ;$$

$$E(a) = 0.2, E(b) = 0.2 ;$$

$$H(a) = 0.3, H(b) = 0.7 ;$$

$$K(a) = 0.3, K(b) = 0.8 .$$

Let  $\tau = \{0, A, 1\}$  and  $\sigma = \{0, H, 1\}$ . Then,

(i)  $B$  is fuzzy pg-closed but not fuzzy preclosed set in  $(X, \tau)$ .

(ii)  $E$  is fuzzy pg-closed but not fuzzy g-closed in  $(X, \tau)$  and  $K$  is fuzzy g-closed but not fuzzy pg-closed set in  $(X, \sigma)$ .

**2.4 DEFINITION :** A function  $f : (X, \tau) \rightarrow (Y, \sigma)$  is said to be approximately fuzzy preirresolute (briefly, ap-Fp-irresolute) if,  $pCl(A) \cdot f^{-1}(H)$  whenever  $H$  is a fuzzy preopen subset of  $(Y, \sigma)$ ,  $A$  is a fuzzy pg-closed subset of  $(X, \tau)$  and  $A \cdot f^{-1}(H)$ .

**2.5 DEFINITION :** A function  $f : (X, \tau) \rightarrow (Y, \sigma)$  is said to be approximately fuzzy preclosed (briefly, ap-Fp-closed) if,  $f(B) \cdot pInt(A)$  whenever  $A$  is a fuzzy pg-closed subset of  $(Y, \sigma)$ ,  $B$  is a fuzzy preclosed subset of  $(X, \tau)$  and  $f(B) \cdot A$ .

**2.6 THEOREM :** (i) A function  $f : (X, \tau) \rightarrow (Y, \sigma)$  is an ap-Fp-irresolute if,  $f^{-1}(A)$  is fuzzy preclosed in  $(X, \tau)$  for every  $A \in FPO(Y)$ .

(ii)  $f : (X, \tau) \rightarrow (Y, \sigma)$  is ap-Fp-closed if,  $f(B) \in FPO(Y)$  for every fuzzy preclosed subset  $B$  of  $(X, \tau)$ .

**PROOF :** (i) Let  $H \cdot f^{-1}(A)$  where  $A \in FPO(Y)$  and  $H$  is a fuzzy pg-closed subset of  $(X, \tau)$ . Therefore  $pCl(H) \cdot pCl(f^{-1}(A)) = f^{-1}(A)$ . Thus,  $f$  is ap-Fp-irresolute.

(ii) Let  $f(B) \cdot A$ , where  $B$  is a fuzzy preclosed subset of  $(X, \tau)$  and  $A$  a fuzzy pg-open subset of  $(Y, \sigma)$ . Therefore,  $pInt(f(B)) \cdot pInt(A)$ . Then,  $f(B) \cdot pInt(A)$ . Thus,  $f$  is ap-Fp-closed.

Clearly, fuzzy preirresolute functions are ap-Fp-irresolute. Also, fuzzy M-preclosed functions are ap-Fp-closed. The converse implication do not hold as it is shown in the following example.

**2.7 EXAMPLE :** Let  $X = \{a, b\}$  and  $Y = \{x, y\}$ . Fuzzy sets  $E, A, B$  and  $H$  be defined as :

$$\begin{aligned} E(a) &= 0, E(b) = 0.8; \\ A(a) &= 0.3, A(b) = 0.4; \\ H(x) &= 0.3, H(y) = 0.4; \\ B(x) &= 0, B(y) = 0.8. \end{aligned}$$

Let  $\tau = \{0, 1\}$  and  $\sigma = \{0, 1\}$ ,  $\rho = \{0, 1\}$  and  $\nu = \{0, 1\}$ . Then, the function  $f : (X, \tau) \rightarrow (Y, \sigma)$  defined by  $f(a) = x$ ,  $f(b) = y$  is ap-Fp-irresolute but not fuzzy preirresolute since  $Z(x) = 0.1$  and  $Z(y) = 0.1$  are fuzzy preopen in  $Y$  but  $f^{-1}(Z)$  is not fuzzy preopen in  $(X, \tau)$ , and the function  $h : (X, \rho) \rightarrow (Y, \nu)$  defined by  $h(a) = x$ ,  $h(b) = y$ , is ap-Fp-closed but not fuzzy M-preclosed.

2.8 REMARK : Converse of the Theorem 2.6 do not hold for .

2.9 EXAMPLE : The function  $f$  defined in Example 2.3 is ap-Fp-irresolute. Fuzzy set  $K(x) = 0.9, K(y) = 0.9$  are fuzzy preopen in  $Y$  but  $f^{-1}(K)$  is not fuzzy preclosed in  $(X, \tau)$ . The function  $h$  defined in Example 2.7 is ap-Fp-closed and the fuzzy set  $W(a) = 0.1, W(b) = 0.1$  is fuzzy preclosed set in  $(X, \tau)$ , but  $h(W)$  is not fuzzy preopen set in  $(Y, \nu)$ .

In the following theorem, we get under certain conditions the converse of the Theorem 2.6 is true.

2.10 THEOREM : Let  $f : (X, \tau) \rightarrow (Y, \sigma)$  be a function. Then,

(i) Let all fuzzy subset of  $(X, \tau)$  be fuzzy clopen, then  $f$  is ap-Fp-irresolute if and only if,  $f^{-1}(H)$  is fuzzy preclosed in  $(X, \tau)$  for every  $H \in FPO(Y)$ .

(ii) Let all fuzzy subsets of  $(Y, \sigma)$  be fuzzy clopen, then  $f$  is ap-Fp-closed if and only if,  $f(B) \in FPO(Y)$  for every fuzzy preclosed subset  $B$  of  $(X, \tau)$ .

PROOF : (i) Assume  $f$  is ap-Fp-irresolute. Let  $A$  be an arbitrary fuzzy subset of  $(X, \tau)$  such that  $A \in Q$  where  $Q \in FPO(X)$ . Then by hypothesis  $pCl(A) \cap pCl(Q) = Q$ . Therefore all fuzzy subsets of  $(X, \tau)$  are fuzzy pg-closed (and hence all are fuzzy pg-open). So for any  $H \in FPO(Y)$ ,  $f^{-1}(H)$  is fuzzy pg-closed in  $X$ . Since  $f$  is ap-Fp-irresolute,  $pCl(f^{-1}(H)) \cap f^{-1}(H) = f^{-1}(H)$ . Therefore,  $pCl(f^{-1}(H)) = f^{-1}(H)$ , i.e.,  $f^{-1}(H)$  is fuzzy preclosed in  $(X, \tau)$ .

The converse is clear by Theorem 2.6.

(ii) Assume  $f$  is ap-Fp-closed. Reasoning as in (i), we obtain that all fuzzy subsets of  $(Y, \sigma)$  are fuzzy pg-open. Therefore, for any fuzzy preclosed subset  $B$  of  $(X, \tau)$ ,  $f(B)$  is fuzzy pg-open in  $Y$ . Since  $f$  is ap-Fp-closed,  $f(B) \cap pInt(f(B)) = pInt(f(B))$ . Therefore,  $f(B) = pInt(f(B))$ , i.e.,  $f(B)$  is fuzzy preopen.

The converse is clear by Theorem 2.6.

As an immediate consequence of Theorem 2.10, we have the following.

(ii)

2.11 COROLLARY : Let  $f : (X, \tau) \rightarrow (Y, \sigma)$  be a function. Then,

(i) Let all the fuzzy subsets of  $(X, \tau)$  be fuzzy clopen, then  $f$  is ap-Fp-irresolute if and only if  $f$  is fuzzy preirresolute.

(ii) Let all the fuzzy subsets of  $(Y, \sigma)$  be fuzzy clopen, the  $f$  is ap-Fp-closed if and only if  $f$  is fuzzy M-preclosed.

### 3. CONTRA FUZZY PRE-IRRESOLUTE FUNCTIONS

**3.1 DEFINITION :** A function  $f : (X, \tau) \rightarrow (Y, \sigma)$  is called :

(i) contra fuzzy preirresolute if  $f^{-1}(A)$  is fuzzy preclosed in  $(X, \tau)$  for each  $A \in FPO(Y)$ ,

(ii) contra fuzzy M-preclosed if  $f(B) \in FPO(Y)$  for each fuzzy preclosed set  $B$  of  $(X, \tau)$ .

**3.2 REMARK :** The concepts of contra fuzzy preirresoluteness and fuzzy preirresoluteness are independent notions. For,

**3.3 EXAMPLE :** Let  $X = \{a, b\}$ ,  $Y = \{x, y\}$  and  $Z = \{p, q\}$ . Define fuzzy sets  $A, E$  and  $B$  as :

$$A(a) = 0.7, A(b) = 0.8;$$

$$E(x) = 0.4, E(y) = 0.3;$$

$$B(p) = 0.3, B(q) = 0.3.$$

Let  $\tau = \{0, A, 1\}$ ,  $\eta = \{0, E, 1\}$  and  $\sigma = \{0, B, 1\}$ . Then the function  $f : (X, \tau) \rightarrow (Y, \eta)$  defined by  $f(a) = x$ ,  $f(b) = y$  is contra fuzzy preirresolute but not fuzzy preirresolute and the function  $h : (Y, \eta) \rightarrow (Z, \sigma)$  defined by  $h(x) = p$  and  $h(y) = q$  is fuzzy preirresolute but not contra fuzzy preirresolute.

In this same manner one can prove that, contra fuzzy M-preclosed functions and fuzzy M-preclosed functions are independent notions.

**3.4 DEFINITION :** A function  $f : (X, \tau) \rightarrow (Y, \sigma)$  is called contra fuzzy precontinuous if  $f^{-1}(A)$  is fuzzy preclosed set in  $(X, \tau)$  for each fuzzy open set  $A$  of  $(Y, \sigma)$ .

**3.5 REMARK :** Every contra fuzzy preirresolute function is contra fuzzy precontinuous, but not conversely. For,

**3.6 EXAMPLE :** Let  $X = \{a, b\}$  and  $Y = \{x, y\}$ . Fuzzy sets  $A$  and  $B$  are defined as :

$$A(a) = 0.7, A(b) = 0.8 ;$$

$$B(x) = 0, B(y) = 0.6 .$$

Let  $\tau = \{0, 1\}$  and  $\sigma = \{0, 1\}$ . Then the function  $f : (X, \tau) \rightarrow (Y, \sigma)$  defined by  $f(a) = x, f(b) = y$  is contra fuzzy precontinuous but not contra fuzzy preirresolute.

The following result can be easily verified. Its proof is straightforward.

**3.7 THEOREM :** Let  $f : (X, \tau) \rightarrow (Y, \sigma)$  be a function. Then the following conditions are equivalent :

- (i)  $f$  is contra fuzzy preirresolute,
- (ii) The inverse image of each fuzzy preclosed set in  $Y$  is fuzzy preopen in  $X$ .

**3.8 REMARK :** By Theorem 2.6, we have that every contra fuzzy preirresolute function is ap-Fp-irresolute and every contra fuzzy preclosed function is ap-Fp-closed, the converse implication do not hold (see Remark 2.8)


Now, we define the following.

**3.9 DEFINITION :** A function  $f : (X, \tau) \rightarrow (Y, \sigma)$  is called perfectly contra fuzzy preirresolute if the inverse image of every fuzzy preopen set in  $Y$  is fuzzy pre-clopen in  $X$

Every perfectly contra fuzzy preirresolute function is contra fuzzy preirresolute and fuzzy preirresolute. But the converse may not be true. For,

**3.10 EXAMPLE :** The functions  $f$  and  $h$  defined in Example 3.3 are contra fuzzy preirresolute and fuzzy preirresolute respectively but not perfectly contra fuzzy preirresolute.

Clearly, the following diagram holds and none of its implications is reversible.



Where contra f.p.i = contra fuzzy preirresolute, ap-Fp-i.= ap-Fp-irresolute, perfectly contra f.p.i.= perfectly contra fuzzy preirresolute.

The following theorem is a decomposition of perfectly contra fuzzy preirresoluteness.

**3.11 THEOREM :** For a function  $f : (X, \tau) \rightarrow (Y, \sigma)$  the following conditions are equivalent:

- (i)  $f$  is perfectly contra fuzzy preirresolute.
- (ii)  $f$  is both contra fuzzy preirresolute and fuzzy preirresolute.

**3.12 THEOREM :** If a function  $f : (X, \tau) \rightarrow (Y, \sigma)$  is ap-Fp-irresolute and fuzzy M-preclosed, then for every Fpg-closed set  $A$  of  $(X, \tau)$ ,  $f(A)$  is Fpg-closed in  $(Y, \sigma)$ .

**PROOF:** Let  $A$  be a Fpg-closed subset of  $(X, \tau)$ . Let  $f(A) \cap B$  where  $B \in FPO(Y)$ . Then  $A \cap f^{-1}(B)$  holds. Since  $f$  is ap-Fp-irresolute  $pCl(A) \cap f^{-1}(B)$  and hence  $f(pCl(A)) \cap B$ . Therefore, we have,  $pCl(f(A)) \cap pCl(f(pCl(A))) = f(pCl(A)) \cap B$ . Hence  $f(A)$  is fuzzy pg-closed in  $(Y, \sigma)$ .

However the following theorem holds. The easy proof of it is omitted.

**3.13 THEOREM :** A function  $f : (X, \tau) \rightarrow (Y, \sigma)$  and  $g : (Y, \sigma) \rightarrow (Z, \gamma)$  be two functions such that  $g \circ f : (X, \tau) \rightarrow (Z, \gamma)$ . Then,

- (i)  $g \circ f$  is contra fuzzy preirresolute, if  $g$  is fuzzy preirresolute and  $f$  is contra fuzzy preirresolute.
- (ii)  $g \circ f$  is contra fuzzy preirresolute, if  $g$  is contra fuzzy preirresolute and  $f$  is fuzzy preirresolute.

In an analogous way, we have the following.

**3.14 THEOREM :** Let function  $f : (X, \tau) \rightarrow (Y, \sigma)$  and  $g : (Y, \sigma) \rightarrow (Z, \gamma)$  be two functions such that  $g \circ f : (X, \tau) \rightarrow (Z, \gamma)$ . Then,

- (i)  $g \circ f$  is ap-Fp-closed, if  $f$  is fuzzy M-preclosed and  $g$  is ap-Fp-closed.
- (ii)  $g \circ f$  is ap-Fp-closed, if  $f$  is ap-Fp-closed and  $g$  is fuzzy M-preopen and  $g^{-1}$  preserves Fpg-closed sets.
- (iii)  $g \circ f$  is ap-Fp-irresolute, if  $f$  is ap-Fp-irresolute and  $g$  is fuzzy preirresolute.

**PROOF :** (i) Suppose  $B$  is an arbitrary fuzzy preclosed subset in  $(X, \tau)$  and  $A$  is a Fpg-open subset of  $(Z, \gamma)$  for which  $(g \circ f)(B) \cap A$ . Then  $f(B)$

is fuzzy preclosed in  $(Y, \sigma)$  because  $f$  is fuzzy M-preclosed. Since  $g$  is ap-Fp-closed,  $g(f(B)) \cdot pInt(A)$ . This implies that  $gof$  is ap-Fp-closed.

(ii) Suppose  $B$  is an arbitrary fuzzy preclosed subset of  $(X, \tau)$  and  $A$  is a fuzzy preopen subset of  $(Z, \gamma)$  for which  $(gof)(B) \cdot A$ . Hence  $f(B) \cdot gi^{-1}(A)$ . Then  $f(B) \cdot pInt(gi^{-1}(A))$  since  $gi^{-1}(A)$  is Fpg-open and  $f$  is ap-Fp-closed. Thus,  $(gof)(B) = g(f(B)) \cdot g(pInt(gi^{-1}(A))) \cdot pInt(ggi^{-1}(A)) \cdot pInt(A)$ . This shows that  $gof$  is ap-Fp-closed.

(iii) Suppose  $H$  is an arbitrary Fpg-closed subset of  $(X, \tau)$  and  $E \in FPO(Z)$  for which  $H \cdot (gof)^{-1}(E)$ . Then  $gi^{-1}(E) \in FPO(Y)$  since  $g$  is fuzzy preirresolute. Since  $f$  is ap-Fp-irresolute,  $pCl(A) \cdot fi^{-1}(gi^{-1}(E)) = (gof)^{-1}(E)$ . This proved that  $gof$  is ap-Fp-irresolute.

In the following result we offer a new characterization of the class of fuzzy pre- $T_{1/2}$  spaces by using the concepts of ap-Fp-irresolute functions and ap-Fp-closed functions.

**3.15 DEFINITION :** A fuzzy topological space  $(X, \tau)$  is said to be fuzzy pre- $T_{1/2}$  space if every Fpg-closed set is fuzzy preclosed in it.

**3.16 THEOREM :** Let  $(X, \tau)$  be a fuzzy topological space. Then the following statements are equivalent:

- (i)  $(X, \tau)$  is fuzzy pre- $T_{1/2}$  space,
- (ii) For every space  $(Y, \sigma)$  and every function  $f : (X, \tau) \rightarrow (Y, \sigma)$ ,  $f$  is ap-Fp-irresolute.

**PROOF:** (i)  $\Rightarrow$  (ii). Let  $E$  be a Fpg-closed subset of  $(X, \tau)$  and suppose that  $E \cdot fi^{-1}(H)$  where  $H \in FPO(Y)$ . Since  $(X, \tau)$  is fuzzy pre- $T_{1/2}$  space,  $E$  is fuzzy preclosed (i.e.,  $E = pCl(E)$ ). Therefore,  $pCl(E) \cdot fi^{-1}(H)$ . Then  $f$  is ap-Fp-irresolute.

(ii)  $\Rightarrow$  (i). Let  $B$  be a Fpg-closed subset of  $(X, \tau)$  and let  $Y$  be the fuzzy set of  $X$  with the topology  $\sigma = 0, B, 1$ . Finally, let  $f : (X, \tau) \rightarrow (Y, \sigma)$  be the identity function. By assumption  $f$  is ap-Fp-irresolute. Since  $B$  is Fpg-closed in  $(X, \tau)$  and fuzzy preopen in  $(Y, \sigma)$  and  $B \cdot fi^{-1}(B)$ , it follows that  $pCl(B) \cdot fi^{-1}(B) = B$ . Hence  $B$  is fuzzy preclosed in  $(X, \tau)$  and therefore,  $(X, \tau)$  is a fuzzy pre- $T_{1/2}$  space.

**3.17 THEOREM :** Let  $(Y, \sigma)$  be a fuzzy topological space. Then the following statements are equivalent :

- (i)  $(Y, \sigma)$  is a fuzzy pre- $T_{1/2}$  space,  
(ii) For every space  $(X, \tau)$  and every function  $f : (X, \tau) \rightarrow (Y, \sigma)$ ,  $f$  is ap-Fp-closed.

Proof : Analogous to Theorem 3.16 making the obvious changes.

### References

- [1] Abd El-Hakeim K. M., Zeyada F. M. and Abd Allah M. A., Fuzzy completely weakly pre-irresolute functions, Bull. Cal. Math. Soc., 93 (2), pp. 121-128, (2001).
- [2] Azad K. K., On fuzzy semicontinuity, fuzzy almost continuity and weakly continuity, J. Math. Anal. Appl., 82, pp. 14-32, (1981).
- [3] Balsubramanian G., On fuzzy pre-separation axioms, Bull. Cal. Math. Soc., 90, pp. 427-434, (1998).
- [4] Bin Shahna A. S., On fuzzy strong semicontinuity and fuzzy pre-continuity, Fuzzy Sets and Systems, 44, pp. 303-308, (1991).
- [5] Bin Shahna A. S., Mapping in fuzzy topological spaces, Fuzzy Sets and Systems, 61, pp. 209-213, (1994).
- [6] Fukutake T., Saraf R. K., Caldas and Mishra S., Mappings via Fgp-closed sets, Bull. Fukuoka Univ. Edu., 52 (III), pp. 11-20, (2003).
- [7] Krstaeska B., Fuzzy strongly preopen sets and fuzzy strongly pre-continuity, Math. Vesnik, 50, pp. 111-123, (1998).
- [8] Park J. H. and Park B. H., Fuzzy pre-irresolute mappings, Pusan Kyongnam Math. J., 10, pp. 303-313, (1995).
- [9] Saraf R. K., On fuzzy contra -preclosed mappings(submitted).
- [10] Singal M. K. and Prakash N., Fuzzy preopen sets and fuzzy pre-separation axioms, Fuzzy Sets and Systems, 44, pp. 273-281, (1991).

- [11] Singh S., Generalization of certain fuzzy topological concepts, Ph. D. Dissertation (1999), R. D. University, Jabalpur (M. P.).
- [12] Zhong B. S., Fuzzy strongly preopen sets and fuzzy strongly pre-continuity, Fuzzy Sets and Systems, 52, pp. 345-351, (1992).

M. Caldas  
Departamento de Matematica Aplicada  
Universidade Federal Fluminense  
Rua Mario Santos Braga, s/n,  
CEP: 24020-140,  
Niteroi, RJ BRASIL  
BRASIL  
e-mail : gmamccs@vm.u®.br

G. B. Navalagi  
Department of Mathematics  
KLE Society's  
G. H. College, Haveri-581110,  
Karnataka  
INDIA  
e-mail : gnavalagi@hotmail.com

R. K. Saraf  
Department of Mathematics  
Govt. K. N. G. College  
Post box 260  
Damoh - 470661 (M. P.)  
INDIA