

Proyecciones Journal of Mathematics
Vol. 31, N° 2, pp. 165-168, June 2012.
Universidad Católica del Norte
Antofagasta - Chile

Hochschild-Serre Statement for the total cohomology

FRANÇOIS LESCURE

UNIVERSITÉ DE LILLE 1, FRANCE

Received : February 2012. Accepted : March 2012

Abstract

Let M be a complex manifold and \mathcal{F} a O_M -module with a \mathfrak{g} -holomorphic action where \mathfrak{g} is a complex Lie algebra (cf. [3]). We denote by $\mathbf{H}(\mathfrak{g}, \mathcal{F})$ the “total cohomology” as defined in [1] [2]. Then we prove that, for any ideal $\mathfrak{a} \subset \mathfrak{g}$, the module $\mathbf{H}^\bullet(\mathfrak{a}, \mathcal{F})$ viewed as a $\mathfrak{g}/\mathfrak{a}$ -module, we have a spectral sequence which converges to $\mathbf{H}(\mathfrak{g}, \mathcal{F})$ and whose E_2 -term is $E_2^{p,q} = H^p(\mathfrak{g}/\mathfrak{a}; \mathbf{H}^q(\mathfrak{a}, \mathcal{F}))$.

Let \mathfrak{g} be a finite dimensional complex Lie algebra and M a complex analytic manifold of finite dimension. Suppose that a holomorphic field \mathbf{u}_M of tangents $(1, 0)$ -vectors on M is associated to each $\mathbf{u} \in \mathfrak{g}$. If this transformation satisfies the condition $[\mathbf{u}_M, \mathbf{v}_M] = [\mathbf{u}, \mathbf{v}]_M$, we shall say that it defines a holomorphic \mathfrak{g} -action on M . To be more precise, the real parts of these fields \mathbf{u}_M are the opposite of the Killing fields of a local holomorphic action of some complex Lie group. Let \mathcal{F} be an O_M -module and, for all $\mathbf{u} \in \mathfrak{g}$, let $\gamma_*(\mathbf{u}) : \mathcal{F} \rightarrow \mathcal{F}$ be a morphism of C -sheaf.

Definition 0.1. *If, for any local section σ of \mathcal{F} and any local holomorphic function f on M , we have:*

- (i) $\gamma_*([\mathbf{u}, \mathbf{v}]) = [\gamma_*(\mathbf{u}), \gamma_*(\mathbf{v})]$
- (ii) $\gamma_*(\mathbf{u})(f\sigma) = \mathbf{L}_{\mathbf{u}_M} f\sigma + f\gamma_*(\mathbf{u})\sigma$,

we say that \mathcal{F} is an O_M -module with a holomorphic \mathfrak{g} -action.

Now, denote by $U(\mathfrak{g}, \mathbf{C})$ be the envelopping algebra of the complex Lie algebra \mathfrak{g} .

In [3], we have introduced the sheaf of crossed algebras $U(\mathfrak{g}, \mathbf{O}_M) \stackrel{\text{def}}{=} \mathbf{O}_M \otimes_{\mathbf{C}} U(\mathfrak{g}, \mathbf{C})$ with the use of the commutation formula: $(1 \otimes \mathbf{u})(\varphi \otimes \mathbf{1}) \stackrel{\text{def}}{=} \mathbf{L}_{\mathbf{u}_M} \varphi \otimes \mathbf{1} + \varphi \otimes \mathbf{u}$. Then, we see immediately that the O_M -modules with a holomorphic \mathfrak{g} -action, are exactly the $U(\mathfrak{g}, \mathbf{O}_M)$ -modules, objects which make some Abelian category denoted $Mod(U(\mathfrak{g}, \mathbf{O}_M))$. On the other hand, in [1] and [2], we have defined, for any holomorphically G -equivariant vector bundle $E \rightarrow M$ (G is a complex Lie group with Lie algebra \mathfrak{g}), the total cohomology denoted $\mathbf{H}^*(\mathfrak{g}, \mathbf{E})$. In [3], we have generalized this total cohomology to any $U(\mathfrak{g}, \mathbf{O}_M)$ -module \mathcal{F} and we have showed indeed that the total cohomology is a derived functor; more precisely, we have proved that:

$$\mathbf{H}^*(\mathfrak{g}, \mathbf{E}) \approx \mathbf{Ext}_{U(\mathfrak{g}, \mathbf{O}_M)}^*(\mathbf{O}_M, \mathbf{E})$$

Proposition 0.2. *Let M , \mathfrak{g} , and so on... be like above. Let \mathcal{F} be a left $U(\mathfrak{g}, \mathbf{O}_M)$ -module and \mathfrak{a} an ideal of the complex Lie algebra \mathfrak{g} . Then:*

- (i) *The total cohomology $\mathbf{H}(\mathfrak{a}, \mathcal{F})$ is naturally a left $(\mathfrak{g}/\mathfrak{a})$ -module.*
- (ii) *There is a Hochschild-Serre spectral sequence E_r whose E_2 -term is given by $H^p(\mathfrak{g}/\mathfrak{a}, \mathbf{H}^q(\mathfrak{a}, \mathcal{F}))$ and which converges to $\mathbf{H}^{p+q}(\mathfrak{g}, \mathcal{F})$*

Proof. (i) It is well known, by the Poincaré-Birkhoff-Witt formula, that $U(\mathfrak{g}, \mathbf{O}_M)$ is a free left $U(\mathfrak{a}, O_M)$ -module, and then also, by the anti-isomorphism T (see [3]), a free right $U(\mathfrak{a}, O_M)$ -module. From this we deduce the exactness of the change of rings functor:

$$U(\mathfrak{g}, \mathbf{O}_M) \otimes_{U(\mathfrak{a}, \mathbf{O}_M)} - : \mathbf{Mod}(U(\mathfrak{a}, \mathbf{O}_M)) \rightarrow \mathbf{Mod}(U(\mathfrak{g}, \mathbf{O}_M))$$

By functor adjunction (see [3]), this exactness allows us to show that the ‘forget functor’: $Mod(U(\mathfrak{g}, \mathbf{O}_M)) \rightarrow \mathbf{Mod}(U(\mathfrak{a}, \mathbf{O}_M))$ preserves injective objects. Also, taking the cohomology of the complex of global \mathfrak{a} -invariant sections of an injective resolution for an $U(\mathfrak{g}, \mathbf{O}_M)$ -module \mathcal{F} , we obtain the total cohomology $\mathbf{H}^\bullet(\mathfrak{a}, \mathcal{F})$ which is then a $(\mathfrak{g}/\mathfrak{a})$ -module and does not depend of the auxiliary choice of the resolution.

(ii) The Grothendieck composition theorem of functors shows that it is sufficient to prove that, if \mathcal{I} is an injective $U(\mathfrak{g}, \mathbf{O}_M)$ -module, then the Chevalley-Eilenberg cohomology $H^p(\mathfrak{g}/\mathfrak{a}, \mathbf{H}^0(\mathfrak{a}, \mathcal{I}))$ of the $(\mathfrak{g}/\mathfrak{a})$ -module $\mathbf{H}^0(\mathfrak{a}, \mathcal{I})$ is zero for $p \geq 1$. For this, we know that it will be enough - and we shall make it - to show that the $\mathbf{H}^0(\mathfrak{a}, \mathcal{I})$ is an injective $(\mathfrak{g}/\mathfrak{a})$ -module.

Indeed, let $0 \rightarrow \mathbf{M}' \xrightarrow{\mathbf{j}} \mathbf{M}$ be a monomorphism of $U(\mathfrak{g}/\mathfrak{a}, \mathbf{C})$ -module. We must factorize each $(\mathfrak{g}/\mathfrak{a})$ -morphism $\mathbf{M}' \xrightarrow{\mathbf{u}} \mathbf{H}^0(\mathfrak{a}, \mathcal{I})$ through the monomorphism \mathbf{j} . Let us consider \mathbf{M}' and \mathbf{M} as \mathfrak{g} -modules with an in-effectiveness \mathfrak{a} ; we introduce, as in [3], the $U(\mathfrak{g}, \mathbf{O}_M)$ -modules $O_{M \otimes_C \mathbf{M}'}$ and $O_{M \otimes_C \mathbf{M}}$, defined by the formula:

$$\gamma_*(\mathbf{u})(\mathbf{f} \otimes \mathbf{m}) = \mathbf{L}_{\mathbf{u}_M} \mathbf{f} \otimes \mathbf{m} + \mathbf{f} \otimes \gamma_*(\mathbf{u})\mathbf{m}.$$

But, \mathbf{j} enlarges it naturally in an arrow of $U(\mathfrak{g}, \mathbf{O}_M)$ -modules $j : O_{M \otimes_C \mathbf{M}'} \rightarrow O_{M \otimes_C \mathbf{M}}$. In more, u allows to define naturally some arrow $O_{M \otimes_C \mathbf{M}'} \rightarrow \mathcal{I}$ which, by the injectivity of \mathcal{I} , factorizes itself by \mathbf{j} with the use of one arrow: $O_{M \otimes_C \mathbf{M}} \rightarrow \mathcal{I}$.

Last arrow that defines one other: $\mathbf{H}^0(\mathfrak{a}, O_{M \otimes_C \mathbf{M}}) \rightarrow \mathbf{H}^0(\mathfrak{a}, \mathcal{I})$. But, then, by restriction of this last arrow to $\mathbf{M} \subset \mathbf{H}^0(\mathfrak{a}, O_{M \otimes_C \mathbf{M}})$, we see easily that this answers the question.

References

- [1] F. Lescure, Action sur la cohomologie de Dolbeault, C. R. Acad. Sci., 314, pp. 923-926, (1992).
- [2] F. Lescure, Cohomologie totale et courants de Dolbeault invariants, J. reine angew. Math., 475, pp. 103-136, (1996).
- [3] F. Lescure La cohomologie totale est un foncteur dérivé, Homology, Homotopy and Applications, Volume 12, Number 1, pp. 367-400, (2010).

François Lescure
U.M.R. CNRS 8524
U.F.R. de Mathématiques
Université de Lille 1
59655 Villeneuve d'Ascq Cedex
FRANCE
e-mail : francois.lescur@math.univ-lille1.fr