



ORIGINAL ARTICLE

Decreased lung compliance increases preload dynamic tests in a pediatric acute lung injury model



Benjamín Erranz^a, Franco Díaz^b, Alejandro Donoso^c, Tatiana Salomón^d,
Cristóbal Carvajal^d, María Fernanda Torres^d, Pablo Cruces^{e,f,*}

^a Centro de Medicina Regenerativa, Facultad de Medicina, Clínica Alemana Universidad del Desarrollo, Santiago, Chile

^b Department of Paediatric, Division of Pediatric Critical Care, University of Alabama at Birmingham, USA

^c Unidad de Paciente Crítico Pediátrico, Hospital Clínico Metropolitano La Florida, Santiago, Chile

^d Unidad de Cuidados Intensivos Pediátrica, Clínica Alemana de Santiago, Chile

^e Unidad de Paciente Crítico Pediátrico, Hospital El Carmen de Maipú, Santiago, Chile

^f Centro de Investigación de Medicina Veterinaria, Escuela de Medicina Veterinaria, Facultad de Ecología y Recursos Naturales, Universidad Andrés Bello, Santiago, Chile

Received 25 February 2015; accepted 9 June 2015

Available online 21 October 2015

KEYWORDS

Hemodynamics;
Preload;
Acute lung injury;
Mechanical
ventilation;
Pediatric

Abstract

Background: Preload dynamic tests, pulse pressure variation (PPV) and stroke volume variation (SVV) have emerged as powerful tools to predict response to fluid administration. The influence of factors other than preload in dynamic preload test is currently poorly understood in pediatrics. The aim of our study was to assess the effect of tidal volume (V_T) on PPV and SVV in the context of normal and reduced lung compliance in a piglet model.

Material and method: Twenty large-white piglets (5.2 ± 0.4 kg) were anesthetized, paralyzed and monitored with pulse contour analysis. PPV and SVV were recorded during mechanical ventilation with a V_T of 6 and 12 mL/kg (low and high V_T , respectively), both before and after tracheal instillation of polysorbate 20.

Results: Before acute lung injury (ALI) induction, modifications of V_T did not significantly change PPV and SVV readings. After ALI, PPV and SVV were significantly greater during ventilation with a high V_T compared to a low V_T (PPV increased from 8.9 ± 1.2 to $12.4 \pm 1.1\%$, and SVV from 8.5 ± 1.0 to $12.7 \pm 1.2\%$, both $P < 0.01$).

Conclusions: This study found that a high V_T and reduced lung compliance due to ALI increase preload dynamic tests, with a greater influence of the latter. In subjects with ALI, lung compliance should be considered when interpreting the preload dynamic tests.

© 2015 Sociedad Chilena de Pediatría. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail address: pcrucesr@gmail.com (P. Cruces).

PALABRAS CLAVE

Hemodinamia;
Precarga;
Lesión Pulmonar
Aguda;
Ventilación Mecánica;
Pediatria

Una compliance pulmonar disminuida incrementa los test dinámicos de precarga en un modelo pediátrico de lesión pulmonar aguda

Resumen

Introducción: Test dinámicos de precarga, variación de presión de pulso (PPV) y variación de volumen sistólico (SVV) han emergido como herramientas poderosas para predecir respuesta a la administración de fluidos. Actualmente la influencia de factores distintos a la precarga en la determinación de los test dinámicos de precarga es pobremente conocida en pediatría. Nuestro objetivo fue medir el efecto del volumen tidal (V_T) sobre PPV y SVV en un contexto de compliance pulmonar normal y disminuida en un modelo porcino.

Material y método: Veinte cerditos Large-White anestesiados y paralizados ($5,2 \pm 0,4$ kg). PPV y SVV fueron medidos por análisis de contorno de pulso durante ventilación con V_T de 6 y 12 mL/kg (V_T bajo y alto, respectivamente), ambos previo y posterior a lesión pulmonar aguda (ALI) químicamente inducida con instilación traqueal de polisorbato 20.

Resultados: Previo a inducción de ALI, PPV y SVV no tuvieron cambios significativos al modificar el V_T . Sin embargo, después de ALI, PPV y SVV fueron significativamente mayores durante ventilación con V_T alto, respecto a V_T bajo (PPV aumentó de $8,9 \pm 1,2$ a $12,4 \pm 1,1\%$, y SVV de $8,5 \pm 1,0$ a $12,7 \pm 1,2\%$, ambos $P < 0,01$).

Conclusiones: Este estudio encontró que un V_T alto y una compliance pulmonar disminuida debido a ALI incrementan los test dinámicos de precarga, con una mayor influencia de esta última. En sujetos con ALI la compliance pulmonar debiera ser considerada al interpretar los test dinámicos de precarga.

© 2015 Sociedad Chilena de Pediatría. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Intravenous fluid administration is a common therapy in critically ill patients, serving as the cornerstone of initial treatment in many conditions. Recent data suggest that fluid management has a major impact on the severity and outcome of critical illness.¹ Knowing that excessive administration of fluids and development of fluid overload are associated with negative effects, the main question when deciding fluid administration is whether there will be beneficial effect on hemodynamics, specifically on cardiac output (CO). Preload dynamic tests, pulse pressure variation (PPV) and stroke volume variation (SVV) have emerged as powerful tools to predict response to fluid administration in different clinical settings.²⁻⁵ Many clinical and experimental studies have determined that PPV >10% and SVV >13% are associated with a significant increase in CO after fluid administration, thus predicting fluid responsiveness. These tests are based on the Frank–Starling Law, which describes the relationship between preload (end-diastolic volume or pressure) and stroke volume. As shown in Fig. 1 two zones can be identified: a steep portion of the curve where changes in preload produce significant changes in CO, and a flat portion of the curve where changes in preload do not significantly change CO (dark gray). Periodic changes in ventricular filling pressures (i.e. preload) occur due to cyclical positive pressure administered during mechanical ventilation. Cyclical changes in stroke volume are greater when the ventricles operate on the steep rather than flat portion of the Frank–Starling curve. Preload dynamic test can identify this particular situation and predict response to fluid administration.⁶ Of note, due to complex cardiopulmonary

interactions, preload dynamic indices do not depend only on cardiac preload.^{7,8} Experimental and small clinical studies have demonstrated that pulmonary and other cardiovascular factors may play a role in the PPV and SVV measurements, though exact mechanisms are not well understood.⁹⁻¹²

Acute lung injury (ALI) is a frequent cause of hypoxemia, loss of respiratory system compliance (C_{RS}) and pulmonary edema. Caution has been claimed respect to the usefulness of dynamic preload tests in patients with ALI where due to low C_{RS} the transmission of positive pressure to the vascular compartment of the lungs, great thoracic veins and the heart may be decreased.¹⁰ On the other hand, subjects with ALI have a decreased compliance resulting in higher changes in tracheal pressure for a given tidal volume (V_T).¹¹ In addition, the standard of care of ALI patients include MV with low V_T . This ventilatory strategy may theoretically decrease the pressure transmitted from the airways to the pleural and pericardial spaces, diminishing the variations of SV and PP.^{12,13} Accordingly, the effect of a decreased V_T on preload dynamic tests in ALI is unpredictable.¹⁴ Because there are marked differences in respiratory and cardiovascular physiology in children with respect to adults (e.g. chest wall compliance, airway wall compliance and resistance, heart rate, stroke volume, pulmonary and systemic vascular resistance, aortic elastance and compliance, metabolic rate), basis for many age-specific differences in the cardiovascular and metabolic responses to injury, these observations should be carefully examined.¹⁵⁻¹⁹

In summary, preload dynamic tests may be an inaccurate measure of filling in cases where C_{RS} or V_T are low, even more so considering the particular features pediatric physiology. The aim of this work was to determine the effect of




Figure 1 Frank–Starling relationship. Once the ventricle is operating near the steep part, volume expansion induces a significant increase in cardiac output. The pulse pressure (PPV) and stroke volume (SVV) variations are marked. By contrast, once the ventricle is functioning on the flat part of the curve, fluid infusion has poor effect on the cardiac output. BP, blood pressure; Paw, airway pressure; PP, pulse pressure.

delivered V_T size on PPV and SVV in normal and reduced lung compliance conditions in a pediatric ALI model.

Material and method

The experimental protocol was approved by Facultad de Medicina Clínica Alemana Universidad del Desarrollo Ethics Committee and the CONICYT (Comisión Nacional de Investigación Científica y Tecnológica) Bioethics Advisory Committee. All of the experimental procedures were consistent with the Guiding Principles in the Care and Use of Laboratory Animals adopted by the American Physiological Society.

Subjects

Twenty anesthetized and mechanically ventilated piglets (5.2 ± 0.4 kg).

Surgical preparation

The trachea was cannulated with a 3.5 mm tracheostomy tube, the left jugular vein with a 4F catheter and the right axillary artery with a 4F thermistor-tipped catheter (PiCCO® PV2014L08, Pulsion Medical Systems, Munich, Germany), all via cut down. Anesthesia and neuromuscular blockade were maintained by continuous infusion of propofol (10 mg/kg/h), fentanyl (4 mcg/kg/h) and pancuronium (0.2 mg/kg/h) during the experiment. Piglets were ventilated in a volume-controlled mode (EVITA XL®/Capnoplus, Dräger Medical, Lübeck, Germany) with a positive end-expiratory pressure of 5 cmH₂O, a V_T of 9 mL/kg, inspiratory time of 0.75 s and a FiO_2 of 0.5. The respiratory rate was adjusted to maintain the end-tidal carbon dioxide concentration (ET-CO₂) at 45 ± 5 mmHg. Normal saline at 5 mL/kg/h was administered during instrumentation. Heart rate, oxygen saturation, central temperature, mean arterial pressure (MAP) and central venous pressure (CVP) were monitored with an Infinity Delta XL® monitor (Dräger Medical, Lübeck,

Germany). Zero pressure was set at the midaxillary line. Temperature was kept at 37.0 ± 0.6 °C.

Acute lung injury

After surgical preparation, each animal was placed in lateral decubitus and a 10% (v/v) solution of polysorbate 20 (Tween® 20 [polyoxyethylene 20 sorbitan monolaurate], Sigma-Aldrich, MO, USA) in saline (1 mL/kg) was instilled in the airway of the dependent lung via a 2-mm catheter. The procedure was repeated with the animal rotated to the opposite side. Residual fluid was suctioned from the airway. Lung injury was targeted to achieve a partial pressure of oxygen (PaO_2) < 200 Torr with $FiO_2 = 1$ in the supine position at 20 min after polysorbate 20 instillation. If this target was not met, polysorbate 20 instillation was repeated:

- (i) *Pulmonary measurements:* Arterial blood gases were determined with an i-STAT® System and i-STAT® Cartridges EG6+ (Abbott Laboratories, Princeton, NJ) from blood samples drawn from the arterial catheter. Oxygenation was assessed by the PaO_2/FiO_2 ratio. Static C_{RS} was calculated as $V_T/(P_{pl} - PEEP_{TOT})$, where P_{pl} is plateau pressure measured after a 4-s inspiratory hold and $PEEP_{TOT}$ is total PEEP measured after a 4-s expiratory hold. These variables were recorded from the ventilator display.
- (ii) *Hemodynamic measurements:* cardiac output (CO) was calculated by transpulmonary thermodilution and continuous CO was obtained with PiCCO system calibrated at baseline and after ALI induction according to manufacturer's instructions. According to studies in humans at rest and under different types of environmental stress, a functional definition of 'normovolemia' would be by its ability to provide the heart with an optimal central blood volume, i.e. that cardiac pumping capacity is not limited by its preload. Functional normovolemia is the point in the cardiac preload/output relationship at which CO does not increase further under circumstances where




Figure 2 Experimental protocol.

venous return is unimpeded.²⁰ Normovolemic state was standardized prior registering hemodynamic data at baseline and after ALI induction, with successive 10 mL/kg intravenous 6% hydroxy-ethyl starch (HES) boluses until CO did not increase by more than 10%. HES was administered over 10 min and personnel deciding fluid administration were blinded for preload dynamic indices.

- (iii) *Preload dynamic indices*: PPV and SVV were obtained from the PiCCO® monitor display. According to PiCCO algorithm SVV is calculated from the mean values of four maximum (SV_{max}) and minimum (SV_{min}) stroke volumes averaged during the previous 30 s (SV_{mean}): $SVV = (SV_{max} - SV_{min}) / SV_{mean}$. PPV is calculated during the same time interval: $PPV = (PP_{max} - PP_{min}) / PP_{mean}$ (2.7). PPV and SVV were registered at baseline and 1 h after ALI induction, with low V_T (6 mL/kg) and high V_T (12 mL/kg). PiCCO® system was calibrated with transpulmonary thermodilution (TPTD) at baseline and after polysorbate 20 instillation and PPV and SVV were registered after 5 min of stability (Fig. 2).

At the end of the study period, while under anesthesia, the animals were euthanized by 10% potassium chloride infusion until the detection of ventricular fibrillation or asystole.

Statistical analysis

Data are expressed in mean \pm SEM. Analysis of variance for repeated measurements was used for comparison between the preload dynamic indicators with high and low V_T . Significance was set at $P < 0.05$.

Results

All subjects completed the experimental protocol. After polysorbate 20 instillation, a significant decrease in PaO_2/FiO_2 from 345 ± 14 to 155 ± 8 Torr and C_{RS} from 1.58 ± 0.13 to 0.93 ± 0.05 mL/cmH₂O/kg was observed (both $P < 0.01$). Also there was an increase in HR (133 ± 7 – 162 ± 8 bpm, $P < 0.001$) without changes in MAP

(74 ± 3 – 76 ± 3 mmHg), CVP (7.6 ± 0.4 – 7.6 ± 0.6 mmHg) and CO (4.1 ± 0.2 – 4.3 ± 0.3 L/min/m²).

Preload dynamic indices

Subjects required 14 ± 2 mL/kg at baseline and 25 ± 3 mL/kg after ALI induction for volemic status optimization. All subjects maintained sinus rhythm during experiment.

At baseline PPV was 8.9 ± 1.2 and $10.7 \pm 1.1\%$ with low and high V_T ($P = 0.058$) and SVV was 8.7 ± 1.2 and $9.7 \pm 1.1\%$ ($P = 0.45$), respectively. After ALI induction PPV increased from 8.9 ± 1.2 to $12.4 \pm 1.1\%$ ($P = 0.037$) and SVV from 8.5 ± 1.0 to $12.7 \pm 1.2\%$ ($P = 0.03$) with low and high V_T respectively (Figs. 3 and 4).

Discussion

The main finding of this experimental study is that high V_T significantly increased PPV and SVV when C_{RS} was reduced, but not when C_{RS} was normal in subjects with optimized intravascular volume (not hypovolemic).

Preload dynamic indices are produced by the transmission of airway pressure to the pleural and pericardial spaces, determining cyclic changes in venous return, cardiac preload and afterload. Therefore, its clinical use could be theoretically limited in conditions where the airway pressure transmitted to the intrathoracic spaces is low (i.e. low V_T and low C_{RS}). Recent studies in anesthetized large animals and critically ill adults reported that V_T size influences preload dynamic indices.^{21,22} Our results differ with previous studies where size of delivered V_T has been an indication as the major determinant of preload dynamic indices.^{10,23} In this model of normovolemic subjects, when V_T was applied in a lung with a reduced C_{RS} the effect was more pronounced, resulting in a leftward shift on the Frank–Starling curve and consequently a greater variation of stroke volume and pulse pressure.²⁴ We found that the reduction of C_{RS} was a factor that amplified the effect of V_T on preload dynamic tests. In a small series of ARDS patients, Vieillard Baron¹³ concluded that V_T and not the airway pressure itself, was the determinant for the




Figure 3 Stroke volume variation (SVV) and pulse pressure variation (PPV) in euvoletic piglets mechanically ventilated with a tidal volume (V_T) 6 and 12 mL/kg at baseline (A) and after ALI induction (B).

afterload and the RV stroke work, both components of the PPV. However, an interesting prior study in a model of oleic acid-lung injury reported that the influence of lung edema in the transmission of airway pressure to the pleural space depends significantly on the peak volume during the inspiration.²⁵ While most studies used oleic acid-lung injury, where the main pathogenic mechanism is an increase in pulmonary vascular permeability, we used a surfactant deactivation model. It has been documented that tracheal instillation of polysorbate 20 produce high-tension pulmonary edema. Despite these differences, the lung edema in subjects with injured lungs was similar to other models of lung injury.^{10,12,16,26} Thus, in models of pulmonary edema, transmission of airway pressure to pleural space and vascular structures may be relevant, and influence preload dynamic markers. Accordingly Huang et al. published a clinical series of patients with severe ARDS ventilated with a low V_T strategy, finding that PPV maintained a good prediction for volume response. Although the V_T reported was approximately 6 mL/kg, the mean peak inspiratory pressure was nearly 34 cmH₂O, obtaining an at least 20 cmH₂O driving pressure.²⁷ These findings agree with our results, since the cyclical changes for the arterial pressure and the volume ejection would be secondary to the generated airway pressure and not to the V_T absolute value.




Figure 4 Changes in stroke volume variation (SVV) and pulse pressure variation (PPV) in each animal mechanically ventilated with a tidal volume (V_T) of 6 and 12 mL/kg after acute lung injury induction.

Additionally we used a model of increased preload (normovolemia as defined, and corroborated by maintaining cardiac output and volume preload after ALI) so that the ventricle is operating on the flat portion of the Frank–Starling curve, not only is the likelihood of finding a change in PPV and SVV with increased V_T reduced, but based on the literature and even in a pediatric model one would not expect the PPV and SVV to change under the condition of V_T of 6 mL/kg. The finding of no increase in PPV and SVV with V_T 12 mL/kg differs from that previously reported^{10,17} and could at least partially be explained by the normovolaemic model used. It would appear that a V_T of 12 mL/kg more closely resembles that of 10 mL/kg rather than 15 mL/kg if this study is compared with that by Renner et al.¹⁷

This study has some limitations. The measurement of pleural pressure would allow us to determine the chest wall compliance and specifically address its effects on preload dynamic markers, but it was not performed due to technical reasons. Additionally, tachycardia and decreased SV after ALI are also known factors that influence preload dynamic markers.

In conclusion, preload dynamic indexes are powerful tools when deciding fluid administration in critically ill subjects, but other factors, such as V_T and C_{RS} , must be

considered for correct interpretation. Future clinical studies should address these variables in addition to hemodynamic parameters and preload dynamic markers for fluid management strategies in critically ill children.

Funding

This work was supported by grant Fondecyt 11075041 from CONICYT (Comisión Nacional de Investigación Científica y Tecnológica, Chile) to Pablo Cruces.

Conflicts of interest

This article meets all requirements about informed consent/acknowledgement, ethics committee, funding, animal research and lack conflict of interest, as appropriate.

References

- Goldstein SL. Fluid management in acute kidney injury. *J Intensive Care Med.* 2014;29:183–9.
- Marik PE, Cavallazzi R, Vasu T, Hirani A. Dynamic changes in arterial waveform derived variables and fluid responsiveness in mechanically ventilated patients: a systematic review of the literature. *Crit Care Med.* 2009;37:2642–7.
- Berkenstadt H, Margalit N, Hadani M, et al. Stroke volume variation as a predictor of fluid responsiveness in patients undergoing brain surgery. *Anesth Analg.* 2001;92:984–9.
- Auler JO, Galas F, Hajjar L, Santos L, Carvalho T, Michard F. Online monitoring of pulse pressure variation to guide fluid therapy after cardiac surgery. *Anesth Analg.* 2008;106:1201–6.
- Cannesson M, Musard H, Desebbe O, et al. The ability of stroke volume variations obtained with Vigileo/FloTrac system to monitor fluid responsiveness in mechanically ventilated patients. *Anesth Analg.* 2009;108:513–7.
- Fujita Y, Yamamoto T, Sano I, Yoshioka N, Hinenoya H. A comparison of changes in cardiac preload variables during graded hypovolemia and hypervolemia in mechanically ventilated dogs. *Anesth Analg.* 2004;99:1780–6.
- Wiesenack C, Prasser C, Rödiger G, Keyl C. Stroke volume variation as an indicator of fluid responsiveness using pulse contour analysis in mechanically ventilated patients. *Anesth Analg.* 2003;96:1254–7.
- Vistisen ST, Koefoed-Nielsen J, Larsson A. Should dynamic parameters for prediction of fluid responsiveness be indexed to the tidal volume? *Acta Anaesthesiol Scand.* 2010;54:191–8.
- Van Lieshout JJ, Harms MP, Pott F, Jenstrup M, Secher NH. Stroke volume of the heart and thoracic fluid content during head-up and head-down tilt in humans. *Acta Anaesthesiol Scand.* 2005;49:1287–92.
- da Silva Ramos FJ, de Oliveira EM, Park M, Schettino GPP, Azevedo LCP. Heart-lung interactions with different ventilatory settings during acute lung injury and hypovolaemia: an experimental study. *Br J Anaesth.* 2011;106:394–402.
- Monnet X, Teboul JL. Pulse pressure variation: does lung compliance really matter. *Crit Care Med.* 2012;40:1691–2.
- Michard F. Changes in arterial pressure during mechanical ventilation. *Anesthesiology.* 2005;103:419–28.
- Vieillard-Baron A, Loubieres Y, Schmitt JM, Page B, Dubourg O, Jardin F. Cyclic changes in right ventricular output impedance during mechanical ventilation. *J Appl Physiol.* 1999;87:1644–50.
- De Backer D, Scolletta S. Why do pulse pressure variations fail to predict the response to fluids in acute respiratory distress syndrome patients ventilated with low tidal volume? *Crit Care.* 2011;15:150.
- Renner J, Meybohm P, Gruenewald M, et al. Global end-diastolic volume during different loading conditions in a pediatric animal model. *Anesth Analg.* 2007;105:1243–9.
- Lopez-Herce J, Ruperez M, Sanchez C, Garcia C, Garcia E. Estimation of the parameters of cardiac function and of blood volume by arterial thermodilution in an infant animal model. *Paediatr Anaesth.* 2006;16:635–40.
- Renner J, Cavus E, Meybohm P, et al. Pulse pressure variation and stroke volume variation during different loading conditions in a paediatric animal model. *Acta Anaesthesiol Scand.* 2008;52:374–80.
- Frey U, Jackson AC, Silverman M. Differences in airway wall compliance as a possible mechanism for wheezing disorders in infants. *Eur Respir J.* 1998;12:136–42.
- Ingimarsson J, Thorsteinsson A, Larsson A, Werner O. Lung and chest wall mechanics in anesthetized children Influence of body position. *Am J Respir Crit Care Med.* 2000;162:412–7.
- Ejlertsen E, Skak C, Moller K, Pott F, Secher NH. Central cardiovascular variables at a maximal mixed venous oxygen saturation in severe hepatic failure. *Transplant Proc.* 1995;27:3506–7.
- Reuter DA, Bayerlein J, Goepfert MS, et al. Influence of tidal volume on left ventricular stroke volume variation measured by pulse contour analysis in mechanically ventilated patients. *Intensive Care Med.* 2003;29:476–80.
- Charron C, Fessenmeyer C, Cosson C, et al. The influence of tidal volume on the dynamic variables of fluid responsiveness in critically ill patients. *Anesth Analg.* 2006;102:1511–7.
- Jardin J, Genevray B, Brun-Ney D, Bourdarias JP. Influence of lung and chest wall compliances on transmission of airway pressure to the pleural space in critically ill patients. *Chest.* 1985;88:653–8.
- Perel A. The physiological basis of arterial pressure variation during positive-pressure ventilation. *Réanimation.* 2005;14:162–71.
- O'Quin RJ, Marini JJ, Culver BH, Butler J. Transmission of airway pressure to pleural space during lung edema and chest wall restriction. *J Appl Physiol.* 1985;59:1171–7.
- Díaz F, Erranz B, Donoso A, et al. Surfactant deactivation in a paediatric model induces hypovolemia and fluid shift to the extravascular lung compartment. *Paediatr Anaesth.* 2013;23:250–7.
- Huang CC, Fu JY, Hu HC, et al. Prediction of fluid responsiveness in acute respiratory distress syndrome patients ventilated with low tidal volume and high positive end-expiratory pressure. *Crit Care Med.* 2008;36:2810–6.